arXiv:math/0312494v3 [math.QA] 4 Dec 2004

Quantum symmetric functions

Rafael Díaz* and Eddy Pariguan†

February 8, 2008

Abstract

We study quantum deformations of Poisson orbivarieties. Given a Poisson manifold $(\mathbb{R}^m, \alpha)$ we consider the Poisson orbivariety $(\mathbb{R}^m)^n/S_n$. The Kontsevich star product on functions on $(\mathbb{R}^m)^n$ induces a star product on functions on $(\mathbb{R}^m)^n/S_n$. We provide explicit formulae for the case $\mathfrak{h} \times \mathfrak{h}/\mathcal{W}$, where $\mathfrak{h}$ is the Cartan subalgebra of a classical Lie algebra $\mathfrak{g}$ and $\mathcal{W}$ is the Weyl group of $\mathfrak{h}$. We approach our problem from a fairly general point of view, introducing Polya functors for categories over non-symmetric Hopf operads.

1 Introduction

Let k be a field of characteristic 0, M be a set and G be a subgroup of the permutation group on n-letters S_n. A function $f : M^n \to k$ is said to be G-symmetric if $f(x_{\sigma(1)}, x_{\sigma(2)}, ..., x_{\sigma(n)}) = f(x_1, x_2, ..., x_n)$ for all $\sigma \in G \subseteq S_n$ and all $x_1, ..., x_n \in M$. The k-space of G-symmetric functions $\mathrm{Func}(M^n, k)^G$ is a subalgebra of the k-algebra $\mathrm{Func}(M^n, k)$ of all functions from M^n to k. One of the goals of this paper is to find explicit formulae for a product on the algebra $\mathrm{Func}(M^n, k)^G$ in a variety of contexts. Our approach is based on the following observations:

- It is often easier to work with coinvariant functions $\mathrm{Func}(M^n, k)_G$ instead of working with invariant functions.
- Symmetric functions arise as an instance of a general construction which assigns to any k-algebra A its n-th symmetric power algebra $\mathrm{Sym}^n(A)$. This insight led us to introduce the notion of Polya functors which we present in the context of categories over non-symmetric Hopf operads.

Our main interest is to study formal deformations of the algebra $\mathrm{Func}(M^n, k)^G$. We take the real numbers $\mathbb{R}$ as the ground field, and let $(\mathbb{R}^m, \{,\})$ be a Poisson manifold. Under this conditions Kontsevich in [14] have shown the existence of a canonical formal deformation $(C^\infty(\mathbb{R}^m)[[\hbar]], \star)$ of the algebra $(C^\infty(\mathbb{R}^m), \cdot)$ of smooth functions on $\mathbb{R}^m$. We prove that if the Poisson bracket on $(\mathbb{R}^m, \{,\})$ is G-equivariant for $G \subset S_m$, then the $\star$-product on $C^\infty(\mathbb{R}^m)[[\hbar]]$ induces a $\star$-product on the algebra of symmetric functions $C^\infty(\mathbb{R}^m)^G[[\hbar]]$, which we call the algebra of *quantum symmetric functions*. We regard this algebra as the deformation quantization of the Poisson orbifold $\mathbb{R}^m/G$. We remark that in a recent paper [6], Dolgushev

---

*Work partially supported by UCV.

†Work partially supported by FONACIT.

has proved the existence of a quantum product on the algebra of invariant functions $C^{\infty}(M)^G[[\hbar]]$ for an arbitrary Poisson manifold M acted upon by a finite group G. His result is based on an alternative proof of the Kontsevich formality theorem which is manifestly covariant.

We present a general description of the quantum product on $\mathbb{R}^m/G$ using the Kontsevich star product. We give explicit formulae for the product rule in the following three cases:

- symplectic orbifold $\mathfrak{h} \times \mathfrak{h}/\mathcal{W}$ where $\mathfrak{h}$ is a Cartan subalgebra of a classical Lie algebra $\mathfrak{g}$, and $\mathcal{W}$ is the Weyl group associated to $\mathfrak{h}$,
- symplectic orbifold $\mathbb{C}^n/\mathbb{Z}_m^n \rtimes S_n$,
- symplectic orbifold $\mathbb{C}^n/\mathcal{D}_m^n \rtimes S_n$, where $\mathcal{D}_m$ is the dihedral group of $2m$ elements.

Our motivation to consider these orbifolds came from the study of noncommutative solitons in orbifolds [9], [17] and the quantization of the moduli space of vacua in M-theory as consider in the matrix model approach. Our results will be raised, to the categorical context in [4]. In a different direction, they may be extended to include the q-Weyl and h-Weyl algebras as it is done in [5]. We would like to mention that these orbifolds have also been studied from a different point of view in [1], and more recently in [8].

We consider the quantum symmetric functions of type A_n and uncover its relation with the $\mathrm{Schur}(\infty, n)$ algebras. The latter algebras are natural generalizations of the Schur algebras as defined in [10]. We also study the symmetric powers of the M-Weyl algebra, which we define as the algebra generated by x^{-1} and $\frac{\partial}{\partial x}$. We provided explicit formulae for the normal coordinates for the M-Weyl algebra as well as for its symmetric powers. Similarly, we make clear the relation between quantum symmetric odd-functions and the Schur algebras of various dimensions . Finally, we give a cohomological interpretation of the algebra of supersymmetric functions.

2 Invariants vs coinvariants

In this section we introduce the notion of Polya functors for categories over non-symmetric Hopf operads, and provide a list of applications of the Polya functors. We will consider invariant theory for finite groups as well as for compact topological groups. To avoid duplication we will consider only the latter case in the proofs.

Let k be a field of characteristic 0 and consider $(\mathbf{Vect}_k, \otimes, k)$ the monoidal category of vector spaces with linear transformations as morphisms. For any set I, consider the category of I-graded vector spaces $\mathbf{Vect}_I$, it has as objects I-graded vector spaces, $V = \bigoplus_{i \in I} V_i, \quad V_i \in \mathrm{Ob}(\mathbf{Vect}_k)$. Morphisms between objects $V, W \in \mathrm{Ob}(\mathbf{Vect}_I)$ are given by $\mathrm{Mor}(V, W) = \prod_{i \in I} \mathrm{Hom}(V_i, W_i)$. The category $(\mathbf{Vect}_I, \otimes_I, k_I)$ has a monoidal structure compatible with direct sums induced by the corresponding structures on $(\mathbf{Vect}_k, \otimes, k)$. Explicitly, given $V, W \in \mathrm{Ob}(\mathbf{Vect}_I)$ we have $(V \oplus W)_i = V_i \oplus W_i$, $(V \otimes W)_i = V_i \otimes W_i$, and $(k_I)_i = k$. For a finite group G, we denote by $\mathbf{Vect}_I(G)$ the category of I-graded vector spaces provided with grading preserving G actions. Morphisms in $\mathbf{Vect}_I(G)$ are intertwiners, i.e., maps $\varphi : V \longrightarrow W$ such that $\varphi(gv) = g\varphi(v)$, for all $v \in V, g \in G$. Abusing notation, for an infinite compact topological group provided with a biinvariant Haar measure $\mathrm{d}g$, we denote by $\mathbf{Vect}_I(G)$ the category of finite dimensional vector spaces over $\mathbb{C}$ provided with a G action. We define the symmetrization map $s_V : V \longrightarrow V^G$ as the map given by

$$\begin{aligned} s_V(v) &= \frac{1}{\mathrm{vol}(G)} \int_G (gv)\mathrm{d}g, \quad \text{if } G \text{ is infinite and } k = \mathbb{C}, \\ s_V(v) &= \frac{1}{\sharp(G)} \sum_{g \in G} gv, \quad \text{if } G \text{ is finite and } k \text{ is a field of characteristic zero,} \end{aligned}$$

where $\sharp(G)$ denotes the cardinality of G. The sequence $0 \longrightarrow \mathrm{Ker}(s_V) \longrightarrow V \longrightarrow V^G \longrightarrow 0$, is exact and we obtain the corresponding commutative triangle

$$\begin{tikzcd} V \arrow[rr] \arrow[dr] & & V^G \\ & V/\mathrm{Ker}(s_V) \arrow[ur, "s_V"'] & \end{tikzcd}$$

We denote the space $V/\mathrm{Ker}(s_V)$ by V_G, and thus we have an isomorphism $s_V : V_G \longrightarrow V^G$. We define two functors

$$\begin{aligned} \mathrm{Inv} : \mathbf{Vect}_I(G) &\longrightarrow \mathbf{Vect}_I \\ V &\longmapsto V^G \end{aligned}$$

$$\begin{aligned} \mathrm{Coinv} : \mathbf{Vect}_I(G) &\longrightarrow \mathbf{Vect}_I \\ V &\longmapsto V_G \end{aligned}$$

For any $v \in V$, $\overline{v}$ denotes the equivalence class of v in V_G. We have the following

Proposition 1. *The maps s_V above define a natural isomorphism $s : \mathrm{Coinv} \longrightarrow \mathrm{Inv}$.*

Proof. For each $V \in \mathbf{Vect}_I(G)$ the construction above provides an isomorphism $s_V : V_G \longrightarrow V^G$. For a given morphism $V \xrightarrow{\alpha} W$, we have

$$\begin{aligned} \alpha \circ s_V(\overline{v}) &= \alpha\left(\frac{1}{\mathrm{vol}(G)} \int_G (gv)\mathrm{d}g\right) = \frac{1}{\mathrm{vol}(G)} \int_G \alpha(gv)\mathrm{d}g \\ &= \frac{1}{\mathrm{vol}(G)} \int_G g(\alpha v)\mathrm{d}g = s_W(\overline{\alpha v}) = s_W \overline{\alpha}(\overline{v}), \quad \text{for all } v \in V \end{aligned}$$

thus proving that for each arrow $V \xrightarrow{\alpha} W$ the diagram

$$\begin{tikzcd} V_G \arrow[r, "s_V"] \arrow[d, "\mathrm{Coinv}(\alpha)"'] & V^G \arrow[d, "\mathrm{Inv}(\alpha)"] \\ W_G \arrow[r, "s_W"] & W^G \end{tikzcd}$$

is commutative. □

Notice that V_G may also be defined as $V_G = V/\langle v - gv : \quad v \in V,\ g \in G\rangle$. Constructions above can be generalized to the category $\mathbf{Cat}_k$ of all k-linear categories. We make a more general construction in the next section in order to include linear categories over non-symmetric Hopf operads.

2.1 Categories over non-symmetric operads

We review the notion of operads and the notion of algebras over operads [15]. For finite groups $G \subset H$ such that G acts on a k-vector space V, the induced representation is defined by $\mathrm{Ind}_G^H(V) = (k[G] \otimes V)_H$ where $k[G]$ denotes the group algebra of G. Let us define two k-linear categories $\mathbb{N}$ and S

- $\mathrm{Ob}(\mathbb{N}) = \{0, 1, 2, \dots, n, \dots\}$
- $\mathrm{Mor}_{\mathbb{N}}(n, m) = \begin{cases} k, & \text{if } m = n \\ 0, & \text{if } m \neq n \end{cases}$
- $\mathrm{Ob}(S) = \{0, 1, 2, \dots, n, \dots\}$
- $\mathrm{Mor}_S(n, m) = \begin{cases} S_n, & \text{if } m = n \\ 0, & \text{if } m \neq n \end{cases}$

The category $\mathrm{Funct}(S^{op}, \mathbf{Vect}_k)$ of contravariant functors from S to $\mathbf{Vect}_k$ possesses three important monoidal structures given on objects by

- $(V + W)(n) = V(n) \oplus W(n)$.
- $V \otimes W(n) = \bigoplus_{i+j=n} \mathrm{Ind}_{S_i \times S_j}^{S_n}(V(i) \otimes_k W(j))$.
- $V \circ W(n) = \bigoplus_{p \geq 0} \bigoplus_{a_1 + \cdots + a_p = n} \mathrm{Ind}_{S_{a_1} \times \cdots \times S_{a_p}}^{S_n} V(p) \otimes_{s_p} W(a_1) \otimes_k \cdots \otimes_k W(a_p)$.

The category $\mathrm{Funct}(\mathbb{N}, \mathbf{Vect}_k)$ admits similar monoidal structures by forgetting the S_n actions.

Definition 2.
- *An operad is a monoid in the monoidal category* $(\mathrm{Funct}(S^{op}, \mathbf{Vect}_k), \circ, 1)$, *where* $1(n) = 0, n \neq 1$ *and* $1(1) = k$.
- *A nonsymmetric operad is a monoid in the monoidal category* $(\mathrm{Funct}(\mathbb{N}, \mathbf{Vect}_k), \circ, 1)$.

Explicitly, an (non-symmetric) operad is given by $m_p : \mathcal{O}(p) \otimes \mathcal{O}(a_1) \otimes \cdots \otimes \mathcal{O}(a_p) \longrightarrow \mathcal{O}(a_1 + \cdots + a_p)$ satisfying the list of axioms given for example in [15]. If no confusion arises we write m instead of m_p.

Definition 3. *Let* $V \in \mathbf{Vect}_k$, *we define the endomorphisms operad by* $\mathrm{End}_V(n) = \mathrm{Hom}(V^{\otimes n}, V)$, *for all* $n \in \mathbb{N}$. *Composition are given by arrows*

$$\begin{tikzcd}
\mathrm{Hom}(V^{\otimes p}, V) \otimes \mathrm{Hom}(V^{\otimes a_1}, V) \otimes \cdots \otimes \mathrm{Hom}(V^{\otimes a_p}, V) \arrow[d] \\
\mathrm{Hom}(V^{\otimes p}, V) \otimes \mathrm{Hom}(V^{\otimes (a_1 + \cdots + a_p)}, V^{\otimes p}) \arrow[d] \\
\mathrm{Hom}(V^{\otimes n}, V)
\end{tikzcd}$$

for integers $n, a_1, \dots, a_p$ *such that* $a_1 + \cdots + a_p = n$. *For more details see* [15].

Let us introduce the category $\mathbf{Pre\text{-}Cat}_k$ of small pre-categories. Objects of $\mathbf{Pre\text{-}Cat}_k$ are called small pre-categories. A small pre-category $\mathcal{C}$ consists of the following data:

- A set of objects $\mathrm{Ob}(\mathcal{C})$.
- A vector space $\mathrm{Mor}_{\mathcal{C}}(x,y)$ associated to each pair of objects $x,y \in \mathrm{Ob}(\mathcal{C})$.

A morphisms $F \in \mathrm{Mor}_{\mathbf{Pre\text{-}Cat}}(\mathcal{C},\mathcal{D})$ from pre-category $\mathcal{C}$ to pre-category D consists of a map $F : \mathrm{Ob}(\mathcal{C}) \longrightarrow \mathrm{Ob}(\mathcal{D})$ and a family of maps $F_{x,y} : \mathrm{Mor}_{\mathcal{C}}(x,y) \longrightarrow \mathrm{Mor}_{\mathcal{D}}(F(x),F(y))$, for $x,y \in \mathrm{Ob}(\mathcal{C})$. $\mathbf{Pre\text{-}Cat}_k$ has a natural partial monoidal structure. Given pre-categories $\mathcal{C}$ and $\mathcal{D}$ such that $\mathrm{Ob}(\mathcal{C}) = \mathrm{Ob}(\mathcal{D}) = X$, we define the product pre-category $\mathcal{C}\mathcal{D}$ as follows

- $\mathrm{Ob}(\mathcal{C}\mathcal{D}) = X$.
- $\mathrm{Mor}_{\mathcal{C}\mathcal{D}}(x,z) = \bigoplus_{y\in X} \mathrm{Mor}_{\mathcal{C}}(x,y) \otimes \mathrm{Mor}_{\mathcal{D}}(y,z)$.

The partial units are the pre-categories k_X, defined as follows

- $\mathrm{Ob}(k_X) = X$.
- $\mathrm{Mor}_{k_X}(a,b) = \begin{cases} k, & \text{if } a = b \\ 0, & \text{otherwise.} \end{cases}$

Given a pre-category $\mathcal{C}$, we define the non-symmetric operad $\mathrm{End}_{\mathcal{C}}(n) = \mathrm{Mor}_{\mathbf{Pre\text{-}Cat}}(\mathcal{C}^n,\mathcal{C})$, $n \in \mathbb{N}$. We used the convention $\mathcal{C}^0 = k_{\mathrm{Ob}(\mathcal{C})}$.

Definition 4. *Let $\mathcal{O}$ be a non-symmetric k-linear operad. An $\mathcal{O}$-category $(\mathcal{C},\gamma)$ is a pre-category $\mathcal{C}$ together with a non-symmetric operad morphism $\gamma : \mathcal{O} \longrightarrow \mathrm{End}_{\mathcal{C}}$. Explicitly, a k-linear $\mathcal{O}$-category $\mathcal{C}$ consist of the following data:*

- *Objects* $\mathrm{Ob}(\mathcal{C})$.
- *Morphisms:* $\mathrm{Hom}_{\mathcal{C}}(x,y) \in \mathrm{Ob}(\mathbf{Vect}_k)$, *for each pair* $x,y \in \mathrm{Ob}(\mathcal{C})$.
- *For each,* $k \in \mathbb{N}$ *and objects* $x_0,x_1,\dots,x_k \in \mathrm{Ob}(\mathcal{C})$ *maps*

$$p_{x_0,\dots,x_k} : \mathcal{O}(k) \otimes \mathrm{Hom}_{\mathcal{C}}(x_0,x_1) \otimes \cdots \otimes \mathrm{Hom}_{\mathcal{C}}(x_{k-1},x_k) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(x_0,x_k)$$

We usually write p instead of $p_{x_0,\dots,x_k}$.

These data should satisfy the following associativity axiom: Given objects $x_0,\dots,x_{n_1+\cdots+n_k}$, and morphisms $a_i \in \mathrm{Hom}_{\mathcal{C}}(x_{i-1},x_i), i \in [n_1+\cdots+n_k]$, $t \in \mathcal{O}(k)$, $t_i \in \mathcal{O}(n_i)$, then

$$p(m(t;t_1,\dots,t_k);a_1,\dots,a_{n_1+\cdots+n_k}) =$$

$$p(t,p(t_1;a_1,\dots,a_{n_1}),\dots,p(t_k;a_{n_1+\cdots+n_{(k-1)}+1},\dots,a_{n_1+\cdots+n_k})).$$

For example if we are given objects $x_i \in \mathrm{Ob}(\mathcal{C})$, for $i = 0, 1, \dots, 5$, morphisms $a_i \in \mathrm{Mor}(x_{i-1}, x_i)$ for $i = 1, \dots, 5$ and $t \in \mathcal{O}(5)$, then the morphism $p(t; a_1, \dots, a_5)$ from object x_0 to object x_5 is represented by the following diagram



Given objects $x_i \in \mathrm{Ob}(\mathcal{C})$, for $i = 0, 1, \dots, 8$, morphisms $a_i \in \mathrm{Mor}(x_{i-1}, x_i)$, $i = 1, 2, \dots, 8$, $t_1, t_3 \in \mathcal{O}(3)$, $t_2 \in \mathcal{O}(2)$ and $t \in \mathcal{O}(4)$, then the axiom from Definition 4 is represented by the following commutative diagram.



Figure 1: Pictorial representation of the associative axiom of Definition 4.

Given a nonsymmetric operad $\mathcal{O}$, we define the category $\mathbf{OCat}_k$ as follows:

- $\mathrm{Ob}(\mathbf{OCat}_k) =$ small $\mathcal{O}$-categories.
- Morphism $\mathrm{Mor}_{\mathcal{O}}(\mathcal{C}, \mathcal{D})$ from $\mathcal{O}$-category $\mathcal{C}$ to $\mathcal{O}$-category $\mathcal{D}$ are functors from $\mathcal{C}$ to $\mathcal{D}$ such that

  $$F(p(t; a_1, \dots, a_n)) = p(t; F(a_1), \dots, F(a_n)),$$

  for given objects $x_0, x_1, \dots, x_n$, morphisms $a_i \in \mathrm{Hom}_{\mathcal{C}}(x_{i-1}, x_1)$, $F(a_i) \in \mathrm{Hom}_{\mathcal{D}}(F(x_{i-1}); F(x_i))$ and $t \in \mathcal{O}(n)$. We call such a functor F an $\mathcal{O}$-functor.

3 Polya Functors

Given an $\mathcal{O}$-category $\mathcal{C}$, $\mathrm{Aut}^1(\mathcal{C}) \subset \mathrm{Funct}_{\mathcal{O}}(\mathcal{C},\mathcal{C})$ denotes the collection of invertible $\mathcal{O}$-functors identical on objects. Let G be a compact topological group. A G-action on an $\mathcal{O}$-category $\mathcal{C}$ is a representation $\rho : G \longrightarrow \mathrm{Aut}^1(\mathcal{C})$. It is defined by a collection of actions $\rho_{x,y} : G \longrightarrow \mathrm{GL}(\mathrm{Hom}_{\mathcal{C}}(x,y))$ such that

$$\rho_{x_0,x_m}(g)(p(t;a_1,\dots,a_m)) = p(t;\rho_{x_0,x_1}(g)(a_1),\dots,\rho_{x_{n-1},x_m}(g)(a_m))$$

for objects $x_0,x_1,\dots,x_n \in \mathrm{Ob}(\mathcal{C})$, $a_i \in \mathrm{Hom}(x_{i-1},x_i)$, for all $i \in [m]$ and $g \in G$. Abusing notation we shall write ga instead of $\rho_{x,y}(g)(a)$ where $x,y \in \mathrm{Ob}(\mathcal{C})$ and $a \in \mathrm{Hom}(x,y)$. We define $\mathbf{OCat}_k(G)$ to be the category of all linear $\mathcal{O}$-categories provided with G actions. Morphisms F from $\mathcal{O}$-category $\mathcal{C}$ to $\mathcal{O}$-category $\mathcal{D}$ are G-equivariant $\mathcal{O}$-functors F from $\mathcal{C}$ into $\mathcal{D}$, i.e., $F(ga) = gF(a)$, for all $a \in \mathrm{Mor}(x,y)$, $g \in G$, where $x,y \in \mathrm{Ob}(\mathcal{C})$. We define

$$\begin{array}{rccc}\mathrm{Inv}: & \mathbf{OCat}_k(G) & \longrightarrow & \mathbf{OCat}_k \\ & \mathcal{C} & \longmapsto & \mathcal{C}^G\end{array}$$

as follows: $\mathrm{Ob}(\mathcal{C}^G) = \mathrm{Ob}(\mathcal{C})$, $\mathrm{Mor}_{\mathcal{C}^G}(x,y) = \mathrm{Mor}_{\mathcal{C}}(x,y)^G$. $\mathcal{C}^G$ is a $\mathcal{O}$-category since

$$gp(t;a_1,\dots,a_m) = p(t;ga_1,\dots,ga_m) = p(t;a_1,\dots,a_m).$$

We define

$$\begin{array}{rccc}\mathrm{Coinv}: & \mathbf{OCat}_k(G) & \longrightarrow & \mathbf{OCat}_k \\ & \mathcal{C} & \longmapsto & \mathcal{C}_G\end{array}$$

as follows: $\mathrm{Ob}(\mathcal{C}_G) = \mathrm{Ob}(\mathcal{C})$, $\mathrm{Mor}_{\mathcal{C}_G}(x,y) = \mathrm{Mor}_{\mathcal{C}}(x,y)_G$ and

$$\overline{p}(t;\overline{a_1},\overline{a_2},\dots,\overline{a_m}) = \frac{1}{\mathrm{vol}(G)^{m-1}}\overline{\int_{G^{(n-1)}} p(t;a_1,g_2a_2,g_3a_3,\dots,g_ma_m)\mathrm{d}g_2\mathrm{d}g_3\dots\mathrm{d}g_m} \tag{1}$$

Theorem 5. *There is a natural isomorphism* $s : \mathrm{Coinv} \longrightarrow \mathrm{Inv}$ *i.e., for* $\mathcal{C} \in \mathrm{Ob}(\mathbf{OCat}_k(G))$ *we are given an isomorphism* $s_{\mathcal{C}} : \mathrm{Coinv}(\mathcal{C}) \longrightarrow \mathrm{Inv}(\mathcal{C})$ *such that* $\mathrm{Inv}(F) \circ s_{\mathcal{C}} = s_{\mathcal{D}} \circ \mathrm{Coinv}(F)$*, for all functors* $F : \mathcal{C} \longrightarrow \mathcal{D}$ *in the category* $\mathbf{OCat}_k(G)$*.*

Proof. Given a category $\mathcal{C}$ and objects $x,y \in \mathrm{Ob}(\mathcal{C})$, let $s_{x,y} : \mathrm{Hom}_{\mathcal{C}}(x,y) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(x,y)^G$ the symmetrization map defined in the Section 2. By Proposition 1, it induces an isomorphism of vector spaces

$$s_{x,y} : \mathrm{Hom}_{\mathcal{C}}(x,y)_G \longrightarrow \mathrm{Hom}_{\mathcal{C}}(x,y)^G.$$

It remains to check that the following diagram is commutative

$$\begin{tikzcd}
 & \mathcal{O}(k) \otimes \mathrm{Hom}_{\mathcal{C}}(x_0,x_1)_G \otimes \cdots \otimes \mathrm{Hom}_{\mathcal{C}}(x_{m-1},x_m)_G \arrow[dl, "p(t,-)"'] \arrow[dr, "\mathrm{id}\otimes s"] & \\
\mathrm{Hom}_{\mathcal{C}}(x_0,x_m)_G \arrow[dr, "s_{x_0,x_m}"'] & & \mathcal{O}(k) \otimes \mathrm{Hom}_{\mathcal{C}}(x_0,x_1)^G \otimes \cdots \otimes \mathrm{Hom}_{\mathcal{C}}(x_{m-1},x_m)^G \arrow[dl, "p(t,-)"] \\
 & \mathrm{Hom}_{\mathcal{C}}(x_0,x_m)^G &
\end{tikzcd}$$

where $s = s_{x_0,x_1} \otimes \cdots \otimes s_{x_{m-1},x_m}$. Clearly,

$$p(t;-)\mathrm{id} \otimes S = \frac{1}{\mathrm{vol}(G)^m} \int_{G^m} p(t; g_1a_1, \dots, g_ma_m)\mathrm{d}g_1\mathrm{d}g_2 \dots \mathrm{d}g_m.$$

On the other hand,

$$\begin{aligned}
s_{x_0,x_m} p(t; \overline{a_1}, \dots, \overline{a_m}) &= \frac{1}{\mathrm{vol}(G)} \int_G g_1 \overline{p(t; a_1, \dots, a_m)} \mathrm{d}g_1 \\
&= \frac{1}{\mathrm{vol}(G)^m} \int_{G^m} g_1 p(t; a_1, h_2a_2, \dots, h_ma_m)\mathrm{d}g_1\mathrm{d}h_2 \dots \mathrm{d}h_m \\
&= \frac{1}{\mathrm{vol}(G)^m} \int_{G^m} p(t; g_1a_1, g_1h_2a_2, \dots, g_1h_ma_m)\mathrm{d}g_1\mathrm{d}h_2 \dots \mathrm{d}h_m \\
&= \frac{1}{\mathrm{vol}(G)^m} \int_{G^m} p(t; g_1a_1, \dots, g_ma_m)\mathrm{d}g_1\mathrm{d}g_2 \dots \mathrm{d}g_m
\end{aligned}$$

making the change of variables $g_2 = g_1h_2, \dots, g_m = g_1h_m$. □

Notice that if $\mathcal{O}$ is an operad then $\mathcal{O} \otimes \mathcal{O}$ is naturally an operad with $(\mathcal{O} \otimes \mathcal{O})(n) = \mathcal{O}(n) \otimes \mathcal{O}(n)$.

Definition 6. *A Hopf operad is an operad together with an operad morphism* $\Delta : \mathcal{O} \longrightarrow \mathcal{O} \otimes \mathcal{O}$.

We have the following lemma

Lemma 7. *If* $\mathcal{O}$ *is a Hopf operad then the category of* $\mathcal{O}$*-algebras is monoidal.*

Proof. If A and B are $\mathcal{O}$-algebras then $A \otimes B$ is also an $\mathcal{O}$-algebra, as the following diagram shows

$$\mathcal{O}(n) \otimes (A \otimes B)^{\otimes n} \longrightarrow (\mathcal{O}(n) \otimes A^{\otimes n}) \otimes (\mathcal{O}(n) \otimes B^{\otimes n}) \longrightarrow A \otimes B.$$

□

We now construct a partial monoidal structure on **OCat**.

Definition 8. *Let* $\mathcal{O}$ *be a Hopf operad, given* $\mathcal{O}$*-categories* $\mathcal{C}$ *and* $\mathcal{D}$ *such that* $\mathrm{Ob}(\mathcal{C}) = \mathrm{Ob}(\mathcal{D}) = X$*, we define the tensor product category* $\mathcal{C} \otimes \mathcal{D}$ *as follows*

- $\mathrm{Ob}(\mathcal{C} \otimes \mathcal{D}) = X$.
- $\mathrm{Mor}_{\mathcal{C}\otimes\mathcal{D}}(x, y) = \mathrm{Mor}_{\mathcal{C}}(x, y) \otimes \mathrm{Mor}_{\mathcal{D}}(x, y)$.
- $p(t; a_1 \otimes b_1, \dots, a_n \otimes b_n) = \sum p(t_{(1)}; a_1, \dots, a_n) \otimes p(t_{(2)}; b_1, \dots, b_n)$, *where* $\Delta(t) = \sum t_{(1)} \otimes t_{(2)}$ *using Swedler notation.*

Recall the well-known definition. Given a pair of groups G and $K \subset S_n$ the semidirect product $G^n \rtimes K$ is the set $G^n \rtimes K = \{(g, a) : g \in G^n, a \in K\}$ provided with the product $(g, a)(h, b) = (ga(h), ab)$ where $g, h \in G^n$, $a, b \in K$ and if $h = (h_1, \dots, h_n)$ then $a(h_1, \dots, h_n) = (h_{a^{-1}(1)}, \dots, h_{a^{-1}(n)})$. The following result is obvious

Lemma 9. *a.* $\mathrm{Ob}(\mathcal{C}^{\otimes n}) = \mathrm{Ob}(\mathcal{C})$, $\mathrm{Mor}_{\mathcal{C}^{\otimes n}}(x,y) = \mathrm{Mor}_{\mathcal{C}}(x,y)^{\otimes n}$.

b. If G acts on $\mathcal{C}$ and $K \subset S_n$ then $G^n \rtimes K$ acts on $\mathcal{C}^{\otimes n}$.

Assume that $\mathcal{C}$ is a category over a non-symmetric Hopf operad $\mathcal{O}$. Let G be a compact topological group, and K a subgroup of S_n. We construct functor $P_{G,K}$ which we call the *Polya functor* of type G, K

$$P_{G,K} : \mathbf{OCat}_k(G) \longrightarrow \mathbf{OCat}_k$$

as follows:

- On objects: Given $\mathcal{C} \in \mathrm{Ob}(\mathbf{OCat}_k(G))$, then $P_{G,K}(\mathcal{C}) \in \mathrm{Ob}(\mathbf{OCat}_k)$ is the category

  $$P_{G,K}(\mathcal{C}) = \mathcal{C}^{\otimes n}/G^n \rtimes K.$$

  Explicitly:

- $\mathrm{Ob}(P_{G,K}(\mathcal{C})) = \mathrm{Ob}(\mathcal{C}^{\otimes n}) = \mathrm{Ob}(\mathcal{C})$, and for given objects $x, y \in \mathrm{Ob}(P_{G,K}(\mathcal{C}))$,

  $$\mathrm{Mor}_{P_{G,K}(\mathcal{C})}(x,y) = (\mathrm{Mor}_{\mathcal{C}}(x,y)^{\otimes n})/G^n \rtimes K.$$

- Identity: $\mathrm{id}_x \in \mathrm{Mor}_{P_{G,K}(\mathcal{C})}(x,x) = \overline{\mathrm{id}_x^{\otimes n}} \in (\mathrm{Mor}_{\mathcal{C}}(x,x)^{\otimes n})/G^n \rtimes K$.

- Composition: Given $x_0, \ldots, x_m \in \mathrm{Ob}(P_{G,K}(\mathcal{C}))$ and morphisms $\overline{a_i} \in \mathrm{Mor}_{P_{G,K}(\mathcal{C})}(x_{i-1}, x_i)$, for $i = 1, \ldots, m$, we have the following

  $$p(t; \overline{a_1}, \ldots, \overline{a_m}) = \frac{1}{((\sharp(G)^n)\sharp(K))^{m-1}} \sum_{(g,s)\in(G^n \rtimes K)^{m-1}} \overline{p(t; a_1, (g_2,s_2)a_2, \ldots, (g_m,s_m)a_m)} \text{ if } G \text{ is finite,}$$

  $$p(t; \overline{a_1}, \ldots, \overline{a_m}) = \frac{1}{(\mathrm{vol}(G^n)\sharp(K))^{m-1}} \sum_{s\in K^{m-1}} \int_{g\in(G^n)^{m-1}} \overline{p(t; a_1, (g_2,s_2)a_2, \ldots, (g_m,s_m)a_m)} \mathrm{d}g_2 \ldots \mathrm{d}g_m,$$

  if G is compact and $g_i \in G^n$.

- On morphisms: each functor $\mathcal{C} \xrightarrow{\alpha} \mathcal{D}$ induces a functor $\mathcal{C}^{\otimes n} \xrightarrow{\alpha^{\otimes n}} \mathcal{D}^{\otimes n}$. This functor descends to a well defined functor

  $$P_{G,K}(\mathcal{C}) = \mathcal{C}^{\otimes n}/G^n \rtimes K \longrightarrow \mathcal{D}^{\otimes n}/G^n \rtimes K = P_{G,K}(\mathcal{D}).$$

Example 10. *Consider the non-symmetric operad $\mathcal{ASS}$ given by $\mathcal{ASS}(n) = k$, for all $n \geq 0$. $\mathcal{ASS}$-categories are categories in the usual sense. This example will be applied in Section 4 to introduce explicit formulae for the composition of morphisms in the Schur categories of various types.*

Definition 11. *Let $\mathcal{O}$ be an operad. An $\mathcal{O}$-algebra is a pair (A, γ) where A is a vector space and $\gamma : \mathcal{O} \longrightarrow \mathrm{End}_A$ is an operad morphism.*

Notice that an $\mathcal{O}$-algebra may be regarded as an $\mathcal{O}$-category $\mathcal{C}$ with one object by setting $A = \mathrm{Mor}_{\mathcal{C}}(1,1)$. We denote by $\mathbf{OAlg}_k$ the category of $\mathcal{O}$-algebras and by $\mathbf{OAlg}_k(G)$ the category of $\mathcal{O}$-algebras provided with a G action. We have two naturally isomorphic functors

$$\begin{array}{rccc} \mathrm{Inv}: & \mathbf{OAlg}_k(G) & \longrightarrow & \mathbf{OAlg}_k \\ & A & \longmapsto & A^G \end{array}$$

$$\begin{array}{rccc} \mathrm{Coinv}: & \mathbf{OAlg}_k(G) & \longrightarrow & \mathbf{OAlg}_k \\ & A & \longmapsto & A_G \end{array}$$

Let $\mathcal{O}$ be a Hopf operad. Assume that A is an $\mathcal{O}$-algebra. Let G be a compact topological group and $K \subset S_n$. We have a functor

$$\begin{array}{rccc} P_{G,K}: & \mathbf{OAlg}_k(G) & \longrightarrow & \mathbf{OAlg}_k \\ & A & \longmapsto & A^{\otimes n}/G^n \rtimes K \end{array}$$

The $\mathcal{O}$-algebra structure on $A^{\otimes n}/G^n \rtimes K$ is given for any $a_1, \ldots, a_m \in A$, $t \in \mathcal{O}(m)$ by

$$p(t; \overline{a_1}, \ldots, \overline{a_m}) = \frac{1}{((\sharp(G)^n)\sharp(K))^{m-1}} \sum_{(g,s)\in(G^n \rtimes K)^{m-1}} \overline{p(t; a_1, (g_2, s_2)a_2, \ldots, (g_m, s_m)a_m)} \tag{2}$$

if G is finite. If G is compact taking $g_i \in G^n$ we have that

$$p(t; \overline{a_1}, \ldots, \overline{a_m}) = \frac{1}{(\mathrm{vol}(G^n)\sharp(K))^{m-1}} \sum_{s\in K^{m-1}} \int_{g\in(G^n)^{m-1}} \overline{p(t; a_1, (g_2, s_2)a_2, \ldots, (g_m, s_m)a_m)} \mathrm{d}g_2 \ldots \mathrm{d}g_m.$$

Corollary 12. *Let A be an $\mathcal{O}$-algebra, G a finite group acting by algebra automorphism on A. Take $K = \{\mathrm{id}\}$. The following identity hold in $P_G(A)$*

$$\overline{a}\ \overline{b} = \frac{1}{\sharp(G)} \sum_{g\in G} \overline{a(gb)} \tag{3}$$

where $a, b \in A$.

We remark that Joyal theory of analytic functors, see [12], [13], may be extended from the context of k-vector spaces to $\mathcal{O}$-algebras by defining a functor F which sends a family $A = \{A_n\}_{n\geq 0}$ of $\mathcal{O}$-algebras provided with right S_n actions, into the functor F_A from the **OAlg** into **OAlg** given as follows

$$\begin{array}{rccc} F: & \mathrm{Funct}(S^{op}, \mathbf{OAlg}) & \longrightarrow & \mathrm{Funct}(\mathbf{OAlg}, \mathbf{OAlg}) \\ & A = \{A_n\}_{n\geq 0} & \longmapsto & F_A(B) = \bigoplus_{n\geq 0} (A_n \otimes B^{\otimes n})_{S_n} \quad \text{for all} \quad B \in \mathrm{Ob}(\mathbf{OAlg}) \end{array}$$

Example 13. *Consider the operad Ass given by $Ass(n) = k[S_n]$, for $n \geq 0$. Ass-algebras are the same as associative algebras. If we take the family $k = \{k\}_n$, $n \geq 0$, provided with the trivial S_n action, then* $F_k(A) = \mathrm{Sym}(A)$

The following remarks justify our choice of name for the Polya functors. Let $K \subset S_n$ be a permutation group. For $k \in K$, let $b_s(k)$ be the number of cycles of k of length s. The *cycle index polynomial* of $K \subset S_n$, is the polynomial in n variables $x_1, \ldots, x_n$

$$P_K(x_1, \ldots, x_n) = \frac{1}{\sharp(K)} \sum_{k\in K} x_1^{b_1(k)} x_2^{b_2(k)} \ldots x_n^{b_n(k)}.$$

Let A be a finite dimensional $\mathcal{O}$-algebra and X a basis for A. Using Polya theory [22], we can compute the dimension of the $\mathcal{O}$-algebra $(A^{\otimes n})_K$ as follows:

$$\begin{aligned}
\dim((A^{\otimes n})_K) &= \sharp(X^n/K) \\
&= P_K(\sharp(X), \sharp(X), \dots, \sharp(X)) \\
&= P_K(\dim A, \dim A, \dots, \dim A).
\end{aligned}$$

3.1 General multiplication rule

For each $K \subset S_n$ consider the Polya functor $P_K : k\text{-alg} \longrightarrow k\text{-alg}$ from the category of associative k-algebras into itself defined on objects as follows: if A is a k-algebra, then $P_K(A)$ denotes the algebra whose underlying vector space is

$$P_K(A) = (A^{\otimes n})/\langle a_1 \otimes \dots \otimes a_n - a_{\sigma^{-1}(1)} \otimes \dots \otimes a_{\sigma^{-1}(n)} : a_i \in A, \sigma \in K \rangle.$$

Our next theorem provides the rule for the product of m elements in $P_K(A)$.

Theorem 14. *For any $a_{ij} \in A$ the following identities holds in $P_K(A)$*

$$\sharp(K)^{m-1} \prod_{i=1}^{m} \left(\overline{\bigotimes_{j=1}^{n} a_{ij}} \right) = \sum_{\sigma \in \{\mathrm{id}\} \times K^{m-1}} \overline{\bigotimes_{j=1}^{n} \left(\prod_{i=1}^{m} a_{i\sigma_i^{-1}(j)} \right)} \tag{4}$$

Proof. It is follows from formula (2), taking Ass as the underlying operad and setting $G = \{\mathrm{id}\}$. □

Theorem 14 implies the following

Proposition 15. *Let A be an algebra provided with a basis $\{e_s | s \in [r]\}$. Assume that $e_s e_t = c(k, s, t) e_k$ (sum over k), for all $s, t \in [r]$. For any given $a = (a_{ij}) \in M_{n \times m}([r])$ the following identity holds in $P_K(A)$*

$$(n!)^{m-1} \prod_{i=1}^{m} \left(\overline{\bigotimes_{j=1}^{n} e_{a_{ij}}} \right) = \sum_{\sigma, \alpha} \left(\prod_{j=1}^{n} c(\alpha^j, a, \sigma) \right) \overline{\bigotimes_{j=1}^{n} e_{\alpha^j_{m-1}}},$$

where the sum runs over all $\sigma \in \{\mathrm{id}\} \times K^{m-1}$, $\alpha = (\alpha_i^j) \in M_{(m-1) \times n}([r])$, and
$c(\alpha^j, a, \sigma) = c(\alpha_1^j, a_{1\sigma_1^{-1}(j)}, a_{2\sigma_2^{-1}(j)}) c(\alpha_2^j, \alpha_1^j, a_{3\sigma_3^{-1}(j)}) \dots c(\alpha_{m-1}^j, \alpha_{m-2}^j, a_{m\sigma_m^{-1}(j)})$.

Proof.

$$\begin{aligned}
(n!)^{m-1} \prod_{i=1}^{m} \left(\overline{\bigotimes_{j=1}^{n} e_{a_{ij}}} \right) &= \sum_{\sigma \in \{\mathrm{id}\} \times G^{m-1}} \overline{\bigotimes_{j=1}^{n} \left(\prod_{i=1}^{m} e_{a_{i\sigma_i^{-1}(j)}} \right)} \\
&= \sum_{\alpha, \sigma} \overline{\bigotimes_{j=1}^{n} c(\alpha, a, \sigma) e_{\alpha_{m-1}}} \\
&= \sum_{\sigma, \alpha} \left(\prod_{j=1}^{n} c(\alpha^j, a, \sigma) \right) \overline{\bigotimes_{j=1}^{n} e_{\alpha^j_{m-1}}}
\end{aligned}$$

□

Let us consider a non-symmetric Hopf operad $\mathcal{O}$. Assume that a basis p_m^t, $t \in [k_m]$, for $\mathcal{O}(m)$ is given for each $m \in \mathbb{N}$. Moreover let us assume that $\Delta(p_m^t) = p_m^t \otimes \cdots \otimes p_m^t$, then the next proposition follows from formula (4).

Proposition 16. *Let A be an $\mathcal{O}$-algebra provided with a basis $\{e_s | s \in [r]\}$, and let $\mathcal{O}(n) = \langle p_m^t : t \in [r_m] \rangle$. Assume that $p_m^t(e_{s_1}, \dots, e_{s_m}) = c(t, k, m, s_1, \dots, s_n)e_k$, (sum over k). For any given $a = (a_{ij}) \in M_{n \times m}([r])$, the following identity holds in $P_K(A)$*

$$p_m^t(\overline{\otimes_{j=1}^n e_{a_{1j}}}, \dots, \overline{\otimes_{j=1}^n e_{a_{mj}}}) = \sum_{\sigma,u} \prod_{j=1}^n c(t, u_j, m, a_{1\sigma_1^{-1}(j)}, \dots, a_{m\sigma_m^{-1}(j)}) \overline{\bigotimes_{j=1}^n e_{u_j}}$$

where the sum runs over all $\sigma \in \{\mathrm{id}\} \times K^{m-1}$ and $u \in [r]^n$.

4 Symmetric power of a supercategory.

Let us consider Polya functor $P_{G,K} : \mathbf{Cat}_k(G) \longrightarrow \mathbf{Cat}_k$ for the case $G = \mathrm{id}$, $K = S_n$, i.e, we consider for each $n \in \mathbb{N}$ the functor

$$\mathrm{Sym}^n : \mathbf{Cat}_k \longrightarrow \mathbf{Cat}_k$$

Recall that a supercategory is a category over the category **Supervect** of $\mathbb{Z}_2$-graded vector spaces with the Koszul rule of signs. Functor Sym^n may be applied to supercategories as well. The next result provides formula for the composition of morphisms in the symmetric powers of a supercategory. We use the notation $a_1 \dots a_n = \overline{a_1 \otimes \cdots \otimes a_n} \in \mathrm{Sym}^n(\mathrm{Hom}_{\mathcal{C}}(x, y))$, for morphisms $a_1, \dots, a_n \in \mathrm{Hom}_{\mathcal{C}}(x, y)$.

Proposition 17. *Let $\mathcal{C}$ be a supercategory and let $a_1, \dots, a_n \in \mathrm{Mor}_{\mathcal{C}}(x, y)$ and $b_1, \dots, b_n \in \mathrm{Mor}_{\mathcal{C}}(y, z)$. In the supercategory $\mathrm{Sym}^n(\mathcal{C})$ the compositions of morphisms is given by*

$$(a_1 a_2 \dots a_n)(b_1 b_2 \dots b_n) = \frac{1}{n!} \sum_{\sigma \in S_n} \mathrm{sgn}(a, b, \sigma)(a_1 b_{\sigma^{-1}(1)})(a_2 b_{\sigma^{-1}(2)}) \dots (a_n b_{\sigma^{-1}(n)})$$

where $\mathrm{sgn}(a, b, \sigma) = (-1)^e$ and $e = e(a, b, \sigma) = \sum_{i>j} \overline{a_i}\,\overline{b_{\sigma^{-1}(j)}} + \sum_{\sigma(i)>\sigma(j)} \overline{b_i}\,\overline{b_j}$.

4.1 Schur categories

Let k be a field of characteristic 0, $m = (m_1, \dots, m_k) \in \mathbb{N}^k$, $\mathbb{Z}_m = \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_k}$ and $n \in \mathbb{N}$. We define the Schur supercategory of type (m, n) as follows:

$$\mathrm{Ob}(S(m, n)) = \text{finite dimesional } k\text{-supervector spaces.}$$

$$\mathrm{Mor}_{S(m,n)}(V, W) = (\mathrm{Hom}_k(V^{\oplus \mathbb{Z}_m}, W^{\oplus \mathbb{Z}_m})^{\otimes n})_{\mathbb{Z}_m^n \rtimes S_n}.$$

$\mathbb{Z}_m^n \rtimes S_n$ acts on $(\mathrm{Hom}_k(V^{\oplus \mathbb{Z}_m}, W^{\oplus \mathbb{Z}_m})^{\otimes n})$ as follows

$$\begin{array}{ccc} \mathbb{Z}_m^n \rtimes S_n \times (\mathrm{Hom}_k(V^{\oplus \mathbb{Z}_m}, W^{\oplus \mathbb{Z}_m})^{\otimes n}) & \longrightarrow & (\mathrm{Hom}_k(V^{\oplus \mathbb{Z}_m}, W^{\oplus \mathbb{Z}_m})^{\otimes n}) \\ ((c_1, \dots, c_n), \sigma)(E_{r_1 s_1}^{t_1 u_1} \dots E_{r_n s_n}^{t_n u_n}) & \longmapsto & (E_{r_{\sigma(1)}(s_1+c_1)}^{t_{\sigma(1)}(u_1+c_1)} \dots E_{r_{\sigma(n)}(s_n+c_n)}^{t_{\sigma(n)}(u_n+c_n)}) \end{array}$$

where $E(V,W)^{kl}_{ij}$ are the elementary linear transformation in $(\mathrm{Hom}_k(V^{\oplus \mathbb{Z}_m}, W^{\oplus \mathbb{Z}_m})$, $i \in [\dim V]$, $k \in [\dim W]$ and $j, l \in \mathbb{Z}_m^k$. We apply Polya functor to obtain explicit formula for the composition rule

$$\mathrm{Mor}(V,W) \otimes \mathrm{Mor}(W,Z) \longrightarrow \mathrm{Mor}(V,Z).$$

Theorem 18. *For any given* $M = m_1 \dots m_k$, $i, t \in [\dim W]$, $k \in [\dim Z]$, $r \in [\dim V]$, *and* $j, l, s, u \in \mathbb{Z}_m^n$, *we have*

$$(E(V,W)^{k_1 l_1}_{i_1 j_1} \dots E(V,W)^{k_n l_n}_{i_n j_n})(E(W,Z)^{t_1 u_1}_{r_1 s_1} \dots E(W,Z)^{t_n u_n}_{r_n s_n}) =$$

$$\frac{1}{M^n n!} \sum_{\substack{\sigma \in S_n \\ t_{\sigma(a)} = i_a}} \mathrm{sgn}(\sigma, i, k, r, t) E(V,Z)^{k_1 l_1}_{r_{\sigma(1)}(s_1 + j_1 - u_1)} \dots E(V,Z)^{k_n l_n}_{r_{\sigma(n)}(s_n + j_n - u_n)}$$

where $\mathrm{sgn}(\sigma, i, k, r, t) = (-1)^e$ *and*

$$e = \sum_{i > j} (i_i + k_i)(r_{\sigma^{-1}(i)} + t_{\sigma^{-1}(i)}) + \sum_{\sigma(i) > \sigma(j)} (r_{\sigma^{-1}(i)} + t_{\sigma^{-1}(i)})(r_{\sigma^{-1}(j)} + t_{\sigma^{-1}(j)}).$$

Proof. Straightforward using Polya functor and Proposition 17. □

We now develop a graphical notation that make transparent the meaning of Theorem 18. Let us assume that $n = 4$, $k = 4$, $m_1 = 6$, $m_2 = 3$, $\dim(V) = 4$, $\dim(W) = 2$ and $\dim(Z) = 3$. We represent an elementary linear transformation $E(V,W)$ as in Figure 2



Figure 2: Representation of elementary transformation.

Notice that each block corresponds to $\mathbb{Z}_6 \times \mathbb{Z}_3$, $\mathbb{Z}_6$ acting horizontally, and $\mathbb{Z}_3$ acting vertically. The number of blocks in the bottom row is $\dim(V)$, and the number of blocks in the top row is $\dim(W)$. Elements of $(\mathrm{Hom}(V^{\oplus \mathbb{Z}_m}, W^{\oplus \mathbb{Z}_m})^{\otimes n})_{(\mathbb{Z}_6 \times \mathbb{Z}_3)^4 \rtimes S_4}$ are depicted by four non-numbered arrows, and similarly for elements of $(\mathrm{Hom}(W^{\oplus \mathbb{Z}_m}, Z^{\oplus \mathbb{Z}_m})^{\otimes n})_{(\mathbb{Z}_6 \times \mathbb{Z}_3)^4 \rtimes S_4}$. Composition is obtained as follows

- Fix an arbitrary enumeration of the arrows in $(\mathrm{Hom}(V^{\oplus \mathbb{Z}_m}, W^{\oplus \mathbb{Z}_m})^{\otimes n})_{(\mathbb{Z}_6 \times \mathbb{Z}_3)^4 \rtimes S_4}$.
- Sum over all possible enumerations of the arrows in $(\mathrm{Hom}(W^{\oplus \mathbb{Z}_m}, Z^{\oplus \mathbb{Z}_m})^{\otimes n})_{(\mathbb{Z}_6 \times \mathbb{Z}_3)^4 \rtimes S_4}$.
- Stacks arrows from $(\mathrm{Hom}(V^{\oplus \mathbb{Z}_m}, W^{\oplus \mathbb{Z}_m})^{\otimes n})_{(\mathbb{Z}_6 \times \mathbb{Z}_3)^4 \rtimes S_4}$ to arrows on $(\mathrm{Hom}(W^{\oplus \mathbb{Z}_m}, Z^{\oplus \mathbb{Z}_m})^{\otimes n})_{(\mathbb{Z}_6 \times \mathbb{Z}_3)^4 \rtimes S_4}$ taking care of enumeration and using the $\mathbb{Z}_2 \times \mathbb{Z}_6$ symmetry.

Notice that composition is interesting in that no-touching arrows may nevertheless be composed (due to the $\mathbb{Z}_6 \times \mathbb{Z}_3$ symmetry) as shown in Figure 3.



Figure 3: Example of composition.

Definition 19. *The Schur superalgebra of type* $(\mathrm{sdimV}, \mathrm{m})$ *is given by* $\mathrm{Hom}_{S_n}(V^{\otimes n}, V^{\otimes n})$*, where* sdim *denotes the superdimension of a supervector space.*

See [10] for more on Schur algebras.

Corollary 20. *For* $m = 1$ *and* $V = W$*,* $\mathrm{Mor}_{S(m,n)}(V, V)$ *is the Schur superalgebra* $\mathrm{Schur}(\mathrm{sdimV}, \mathrm{m})$ *of type* $(\mathrm{sdimV}, \mathrm{m})$.

Proof. $(\mathrm{Hom}(V, V)^{\otimes n})_{S_n} \cong (\mathrm{Hom}(V, V)^{\otimes n})^{S_n} \cong \mathrm{Hom}_{S_n}(V^{\otimes n}, V^{\otimes n})$. □

5 Classical symmetric functions

In this section we study classical symmetric functions by means of the Polya functor. We provide a fairly elementary interpretation of the symmetric functions in terms of the symmetric powers of the monoidal algebra associated to the additive monoid $\mathbb{N}^m$. We also consider symmetric odd-functions as well as symmetric Boolean algebras. Symmetric functions have been studied from many points of view, see for example [16], [19], [21] .

5.1 Symmetric functions of Weyl type

The classical Weyl groups of type A_n, B_n, and D_n, are S_n, $\mathbb{Z}_2^n \rtimes S_n$ and $\mathbb{Z}_2^{n-1} \rtimes S_n$ respectively. These groups act on $(\mathbb{R}^m)^n$ as follows:

$$\begin{array}{ccc} S_n \times (\mathbb{R}^m)^n & \longrightarrow & (\mathbb{R}^m)^n \\ (\sigma, (x_1, \ldots, x_n)) & \longmapsto & (x_{\sigma^{-1}(1)}, \ldots, x_{\sigma^{-1}(n)}) \end{array}$$

$$\begin{array}{ccc} (\mathbb{Z}_2^n \rtimes S_n) \times (\mathbb{R}^m)^n & \longrightarrow & (\mathbb{R}^m)^n \\ ((t_1, \ldots, t_n), \sigma)(x_1, \ldots, x_n) & \longmapsto & (t_1 x_{\sigma^{-1}(1)}, \ldots, t_n x_{\sigma^{-1}(n)}) \end{array}$$

The group D_n is regarded as a subgroup of B_n as follows

$$\mathbb{Z}_2^{n-1} \rtimes S_n = \{((t_1, \ldots, t_n), \sigma) \in \mathbb{Z}_2^n \rtimes S_n : t_1 t_2 \ldots t_n = 1\}.$$

Definition 21. *Fix* $m \in \mathbb{N}$*. The algebra of symmetric functions of type* A_n*,* B_n *and* D_n *are given by*

- $\mathrm{Sym}_{A_n}(m) = (\mathbb{C}[x_1, \ldots, x_n])_{S_n} \cong (\mathbb{C}[x_1, \ldots, x_n])^{S_n}$,
- $\mathrm{Sym}_{B_n}(m) = (\mathbb{C}[x_1, \ldots, x_n])_{\mathbb{Z}_2^n \rtimes S_n} \cong (\mathbb{C}[x_1, \ldots, x_n])^{\mathbb{Z}_2^n \rtimes S_n}$,

- $\mathrm{Sym}_{D_n}(m) = (\mathbb{C}[x_1,\ldots,x_n])_{\mathbb{Z}_2^{n-1}\rtimes S_n} \cong (\mathbb{C}[x_1,\ldots,x_n])^{\mathbb{Z}_2^{n-1}\rtimes S_n}$,

where $x_i = (x_{i1},\ldots,x_{im})$, *for* $i = 1,\ldots,n$.

The map $\mathbb{C}[\mathbb{N}^m]^{\otimes n} \longrightarrow \mathbb{C}[x_1,\ldots,x_n]$ given by $a_1\otimes\cdots\otimes a_n \longmapsto x_1^{a_1}\ldots x_n^{a_n}$ defines an isomorphism of algebras, where $a_i \in \mathbb{N}^m$ and $x_i^{a_i} = x_{i1}^{a_{i1}}\ldots x_{im}^{a_{im}}$. We set $\mathbb{N}_e^m = \{a \in \mathbb{N}^m : |a| \text{ is even}\}$ and $\mathbb{N}_o^m = \{a \in \mathbb{N}^m : |a| \text{ is odd}\}$. We denote $X^A = x_1^{a_1}\ldots x_n^{a_n}$, for $A = (a_1,\ldots,a_n) \in (\mathbb{N}^m)^n$.

Theorem 22 (Classical symmetric functions).

a. *The following is a commutative diagram*

$$\begin{tikzcd}
(\mathbb{C}[\mathbb{N}^m]^{\otimes n})_{S_n} \arrow[r, phantom, "\supset"] \arrow[d] & (\mathbb{C}[\mathbb{N}_e^m + \mathbb{N}_o^m]^{\otimes n})_{S_n} \arrow[r, phantom, "\supset"] \arrow[d] & (\mathbb{C}[\mathbb{N}_e^m]^{\otimes n})_{S_n} \arrow[d] \\
\mathrm{Sym}_{A_n}(m) \arrow[r, two heads] & \mathrm{Sym}_{D_n}(m) \arrow[r, two heads] & \mathrm{Sym}_{B_n}(m)
\end{tikzcd}$$

where the vertical arrows are isomorphisms.

b. *For* $A, B \in (\mathbb{N}^m)^n$, *the product rule in* $\mathrm{Sym}_{A_n}(m)$ *is given by*

$$\overline{X^A}\ \ \overline{X^B} = \frac{1}{n!}\sum_{\sigma\in S_n}\overline{X^{A+\sigma(B)}}$$

and on $\mathrm{Sym}_{B_n}(m)$ *and* $\mathrm{Sym}_{D_n}(m)$ *by restriction.*

Proof. The first row of the diagram above follows from the isomorphism above after taken care of the $\mathbb{Z}_2^n$ (resp. $\mathbb{Z}_2^{n-1}$) symmetries for the groups B_n and D_n respectively. It is clear that part b implies the rest of a. We prove b. Given $A, B \in (\mathbb{N}^m)^n$, using Proposition 17 for the product in $\mathrm{Sym}_{A_n}(m)$, we obtain

$$\begin{aligned}
\overline{X^A}\ \ \overline{X^B} &= \frac{1}{n!}\sum_{\sigma\in S_n}\overline{(x_1^{a_1}\ldots x_n^{a_n})(x_1^{b_{\sigma^{-1}(1)}}\ldots x_n^{b_{\sigma^{-1}(n)}})} \\
&= \frac{1}{n!}\sum_{\sigma\in S_n}\overline{X^{A+\sigma(B)}}.
\end{aligned}$$

Now consider $A, B \in (\mathbb{N}_e^m)^n$, the product in $\mathrm{Sym}_{B_n}(m)$ is given by

$$\begin{aligned}
\overline{X^A}\ \ \overline{X^B} &= \frac{1}{2^n n!}\sum_{(t,\sigma)\in\mathbb{Z}_2^n\rtimes S_n}\overline{x_1^{a_1}\ldots x_n^{a_n}t_1^{b_1}x_1^{b_{\sigma^{-1}(1)}}\ldots t_n^{b_n}x_n^{b_{\sigma^{-1}(n)}}} \\
&= \frac{1}{2^n n!}\sum_{\sigma\in S_n}\left(\sum_{t\in\mathbb{Z}_2^n}(-1)^{\sum_{t_i=-1} b_i}\right)\overline{X^{A+\sigma(B)}} \\
&= \frac{1}{n!}\sum_{\pi\in S_n}\overline{X^{A+\sigma(B)}}, \quad \text{since} \sum_{t\in\mathbb{Z}_2^n}(-1)^{\sum_{t_i=-1} b_i} = 2^n.
\end{aligned}$$

Consider $A, B \in (\mathbb{N}_e^n + \mathbb{N}_o^m)^n$, we obtain

$$\begin{aligned}
\overline{X^A}\ \ \overline{X^B} &= \frac{1}{2^{n-1}n!} \sum_{(t,\sigma)\in\mathbb{Z}_2^n \rtimes S_n} \overline{x_1^{a_1}\dots x_n^{a_n} t_1^{b_1} x_1^{b_{\sigma^{-1}(1)}} \dots t_n^{b_n} x_n^{b_{\sigma^{-1}(n)}}} \\
&= \frac{1}{2^{n-1}n!} \sum_{\pi\in S_n} \left(\sum_{\substack{t\in\mathbb{Z}_2^n \\ \prod t_i=1}} (-1)^{\sum_{t_i=-1} b_i + b_n} \right) \overline{X^{A+\sigma(B)}} \\
&= \frac{1}{n!} \sum_{\sigma\in S_n} \overline{X^{A+\sigma(B)}}, \quad \text{since} \quad \sum_{\substack{t\in\mathbb{Z}_2^n \\ \prod t_i=1}} (-1)^{\sum_{t_i=-1} b_i + b_n} = 2^{n-1}
\end{aligned}$$

□

5.2 Symmetric odd-functions

Consider the alternating algebra $\bigwedge[\theta_1, \dots, \theta_m]$, which we regard as the algebra of functions on the purely odd super-space $\mathbb{R}^{0|m}$. A basis for $\bigwedge[\theta_1, \dots, \theta_m]$ is given by $\{\theta_I = \theta_{i_1} \dots \theta_{i_k} |\ I \subset [m]\}$. The structural coefficients are given by $\theta_I\theta_J = c(I,J)\theta_{I\cup J}$, where $c(I,J) = (-1)^{|\{i\in I, j\in J,\ i>j\}|}$, if $I \cap J = \emptyset$, and 0 otherwise. The algebra $\bigwedge[\theta_1, \dots, \theta_m]$ is $\mathbb{Z}_2$-graded, with grading $\overline{\theta_I} = \overline{I} = 0$ if $|I|$ is even and $\overline{\theta_I} = \overline{I} = 1$ if $|I|$ is odd. Now we apply Polya functor to obtain

Proposition 23. *The product rule in the algebra of symmetric odd functions $(\bigwedge[\theta_1, \dots, \theta_m]^{\otimes n})_{S_n}$ is given by*

$$(\overline{\theta_{I_1} \dots \theta_{I_n}})(\overline{\theta_{J_1} \dots \theta_{J_n}}) = \frac{1}{n!} \sum_{\sigma\in S_n} \left(\operatorname{sgn}(I,J,\sigma) \prod_{k=1}^{n} c(I_k, J_{\sigma^{-1}(k)}) \right) \overline{\prod_{i=1}^{n} \theta_{I_k \cup J_{\sigma^{-1}(k)}}},$$

where $\operatorname{sgn}(I,J,\sigma) = (-1)^e$ *and* $e = \sum_{k>l} \overline{I_k J_{\sigma^{-1}(l)}} + \sum_{\sigma(k)>\sigma(l)} \overline{J_k J_l}$.

Proof.

$$\begin{aligned}
(\overline{\theta_{I_1} \dots \theta_{I_n}})(\overline{\theta_{J_1} \dots \theta_{J_n}}) &= \frac{1}{n!} \sum_{\sigma\in S_n} \operatorname{sgn}(I,J,\sigma)(\theta_{I_1}\theta_{J_{\sigma^{-1}(1)}}) \dots (\theta_{I_n}\theta_{J_{\sigma^{-1}(n)}}) \\
&= \frac{1}{n!} \sum_{\sigma\in S_n} \operatorname{sgn}(I,J,\sigma) \left(\prod_{k=1}^{n} c(I_k, J_{\sigma^{-1}(k)}) \theta_{I_k \cup J_{\sigma^{-1}(k)}} \right) \\
&= \frac{1}{n!} \sum_{\sigma\in S_n} \left(\operatorname{sgn}(I,J,\sigma) \prod_{k=1}^{n} c(I_k, J_{\sigma^{-1}(k)}) \right) \overline{\prod_{k=1}^{n} \theta_{I_k \cup J_{\sigma^{-1}(k)}}}
\end{aligned}$$

□

5.3 Symmetric Boolean algebra

Fix $n \in \mathbb{N}$ and let $P(n)$ be the free $\mathbb{C}$-vector space generated by the subsets of $[n]$, i.e., $P(n) = \langle A : \ A \subset [n] \rangle$. Define a product $\cup$ on $P(n)$ by

$$\begin{array}{rccc} \cup : & P(n) \otimes P(n) & \longrightarrow & P(n) \\ & A \otimes B & \longmapsto & A \cup B \end{array}$$

$(P(n), \cup)$ is a Boolean algebra and $\dim(P(n)) = 2^n$. S_n acts naturally on $[n]$ and thus on $P(n)$. We call the algebra $(P(n), \cup)/S_n \cong (P(1)^{\otimes n})/S_n = \mathrm{Sym}^n(P(1))$ the *symmetric Boolean algebra*; it has dimension $n+1$, a basis being $\{[\overline{0}], [\overline{1}], \ldots, [\overline{n}]\}$. We define $P(n,k) := \{A \subset [n] : |A| = k\}$. An application of Polya functor yields the next

Theorem 24. $[\overline{a}][\overline{b}] = \dfrac{1}{\binom{n}{b}} \displaystyle\sum_{k=0}^{m} \binom{a}{b-k} \binom{n-a}{k} \overline{[a+k]}$, *for all* $[\overline{a}], [\overline{b}] \in (P(n), \cup)_{S_n}$, *and* $m = \min(b, n-a)$.

Proof.

$$\begin{aligned} [\overline{a}][\overline{b}] &= \frac{1}{n!} \sum_{\sigma \in S_n} \overline{[a] \cup \sigma[b]} = \frac{1}{\binom{n}{b}} \sum_{B \in P[n,b]} \overline{[a] \cup B} \\ &= \frac{1}{\binom{n}{b}} \sum_{\substack{B_0 \subset P([n]-[a],k) \\ B_1 \subset P([a], b-k)}} \overline{[a] \cup B_0} = \frac{1}{\binom{n}{b}} \sum_{k=0}^{m} \binom{a}{b-k} \binom{n-a}{k} \overline{[a+k]} \end{aligned}$$

□

6 Quantum symmetric functions

In this section we assume the reader is familiar with the notations from [14]. Let us recall the notion of a formal deformation

Definition 25. *Fix a Poisson manifold* $(M, \{,\})$. *A formal deformation (deformation quantization) of the algebra of smooth functions on* M *is an associative star product* $\star : C^\infty(M)[[\hbar]] \otimes_{\mathbb{R}[[\hbar]]} C^\infty(M)[[\hbar]] \longrightarrow C^\infty(M)[[\hbar]]$ *such that:*

- *a.* $f \star g = \displaystyle\sum_{n=0}^{\infty} B_n(f,g)\hbar^n$, *where* $B_n(-,-)$ *are bi-differential operators .*
- *b.* $f \star g = fg + \frac{1}{2}\{f,g\}\hbar + O(\hbar^2)$, *where* $O(\hbar^2)$ *are terms of order* $\hbar^2$.

In [14] a canonical $\star$-product has been constructed for any Poisson manifold. For manifold $(\mathbb{R}^m, \alpha)$ with Poisson bivector α, the $\star$-product is given by the formula

$$f \star g = \sum_{n=0}^{\infty} \frac{\hbar^n}{n!} \sum_{\Gamma \in G_n} \omega_\Gamma B_{\Gamma,\alpha}(f,g),$$

where G_n is a collection of *admissible graphs* each of which has n edges, and ω_Γ are some constants (independent of the Poisson manifold). Given a finite group K acting on $(C^\infty(M), \star)$ by algebra automorphisms, we call the algebra $(C^\infty(M)[[\hbar]], \star)_K \cong (C^\infty(M)[[\hbar]], \star)^K$ the algebra of *quantum K-symmetric functions* on M.

Next theorem shows how groups of automorphisms of $(C^\infty(M)[[\hbar]], \star)$ arise in a natural way.

Theorem 26. *Assume we are given a Poisson structure* $\{\text{-},\text{-}\}$ *on* $\mathbb{R}^m$*, and a group* $K \subset S_m$ *such that* $\{\text{-},\text{-}\}$ *is* K*-equivariant. Then* K *acts on* $(C^\infty(\mathbb{R}^m)[[\hbar]], \star)$ *by automorphisms.*

Proof. We assume that $\{f, g\} \circ \sigma = \{f \circ \sigma, g \circ \sigma\}$, for all $f, g \in C^\infty(\mathbb{R}^m)$, $\sigma \in K$, or equivalently $\alpha^{ij}(\sigma x) = \alpha^{\sigma(i)\sigma(j)}$, where $\alpha^{ij} = \{x_i, x_j\}$ for all $i, j \in [m]$. Let us show that $\sigma(f \star g) = (\sigma f) \star (\sigma g)$.

$$\sigma(f \star g)(x) = (f \star g)(\sigma^{-1}x) = \sum_{n=0}^{\infty} \frac{\hbar^n}{n!} \sum_{\Gamma} w_\Gamma B_{\Gamma,\alpha}(f, g)(\sigma^{-1}x).$$

On the other hand

$$(\sigma f) \star (\sigma g)(x) = \sum_{n=0}^{\infty} \frac{\hbar^n}{n!} \sum_{\Gamma} w_\Gamma B_{\Gamma,\alpha}(\sigma f, \sigma g)(x).$$

We need to prove that

$$B_{\Gamma,\alpha}(f, g)(\sigma^{-1}x) = B_{\Gamma,\alpha}(\sigma f, \sigma g)(x), \quad \text{for all } \ \Gamma \in G_n.$$

Using Kontsevich's formula (see [14]) we get

$$\begin{aligned}
B_{\Gamma,\alpha}(f, g)(\sigma^{-1}x) \quad &= \quad \sum_{I : E_\Gamma \longrightarrow [m]} \left[\prod_{i=1}^{n} \left(\prod_{e \in E_\Gamma, e=(*,i)} \partial_{I(e)} \right) \alpha^{I(e_i^1) I(e_i^2)} \right] (\sigma^{-1}x) \times \\
&\quad \left(\left(\prod_{e \in E_\Gamma, e=(*,L)} \partial_{I(e)} \right) f \right) (\sigma^{-1}x) \times \left(\left(\prod_{e \in E_\Gamma, e=(*,R)} \partial_{I(e)} \right) g \right) (\sigma^{-1}x) \\
&= \quad \sum_{I : E_\Gamma \longrightarrow [m]} \left[\prod_{i=1}^{n} \left(\prod_{e \in E_\Gamma, e=(*,i)} \partial_{\sigma(I(e))} \right) \alpha^{\sigma(I(e_i^1)) \sigma(I(e_i^2))} \right] (x) \times \\
&\quad \left(\left(\prod_{e \in E_\Gamma, e=(*,L)} \partial_{\sigma(I(e))} \right) f \circ \sigma^{-1} \right) (x) \times \left(\left(\prod_{e \in E_\Gamma, e=(*,R)} \partial_{\sigma(I(e))} \right) g \circ \sigma^{-1} \right) (x) \\
&= \quad B_{\Gamma,\alpha}(\sigma f, \sigma g)(x).
\end{aligned}$$

□

Corollary 27. *Under the conditions above, the product rule on* $(C^\infty((\mathbb{R}^m)^n)[[\hbar]], \star)_K$ *is given by*

$$\overline{f} \star \overline{g} = \sum_{\sigma \in K} \sum_{n=0}^{\infty} \frac{\hbar^n}{n!} \left(\sum_{\Gamma} w_\Gamma \overline{B_{\Gamma,\alpha}(f, g \circ \sigma^{-1})} \right) \tag{5}$$

for all $f, g \in C^\infty(\mathbb{R}^m)^n[[\hbar]]$.

Proof. Using Polya functor corollary 12, we have

$$\overline{f} \star \overline{g} = \sum_{\sigma\in K} \overline{f\star\sigma g} = \sum_{\sigma\in K}\sum_{n=0}^{\infty}\frac{\hbar^n}{n!}\left(\sum_{\Gamma} w_\Gamma \overline{B_{\Gamma,\alpha}(f, g\circ\sigma^{-1})}\right).$$

□

Definition 28. *Given a Poisson manifold* $(\mathbb{R}^m, \{\ ,\ \})$ *and a subgroup* $K \subset S_n$ *the algebra of* quantum symmetric functions *on* $(\mathbb{R}^m)^n$ *is set to be* $(C^\infty(\mathbb{R}^m)^n[[\hbar]], \star)_K \cong (C^\infty(\mathbb{R}^m)^n[[\hbar]], \star)^K$.

Notice that if $(\mathbb{R}^m, \alpha)$ is a Poisson manifold then $(\mathbb{R}^m)^n$ is a Poisson manifold in a natural way. Moreover the Poisson structure on $(\mathbb{R}^m)^n$ is S_n-equivariant, and thus K-equivariant for all subgroup K of S_n.

6.1 Weyl algebra

The Kontsevich $\star$-product given by formula $f\star g = \sum_{n=0}^{\infty}\frac{\hbar^n}{n!}\sum_{\Gamma\in G_n}\omega_\Gamma B_{\Gamma,\alpha}(f,g)$ is notoriously difficult to compute. Nevertheless, there are two main examples, see [14] in which a fairly explicit knowledge of the start product is available:

a. If α is a constant non-degenerated Poisson bracket on $\mathbb{R}^{2n}$, then the quantum algebra of polynomial functions on $\mathbb{R}^{2n}$, i.e., $(\mathbb{C}[x_1,\dots,x_{2n}][[\hbar]], \star)$ is isomorphic to $W \otimes_{\mathbb{C}[[\hbar]]} \cdots \otimes_{\mathbb{C}[[\hbar]]} W$, where W is the Weyl algebra, (see definition below).

b. If α is linear Poisson bracket in $\mathbb{R}^n$, then $(\mathbb{R}^n, \alpha)$ is isomorphic as a Poisson manifold to $\mathfrak{g}^*$ for some Lie algebra $\mathfrak{g}$. In this case the quantum algebra of polynomial functions on $\mathfrak{g}^*$, i.e., $(\mathbb{C}[\mathfrak{g}^*][[\hbar]], \star)$ is isomorphic to the universal enveloping algebra $U_h(\mathfrak{g})$ of $\mathfrak{g}$.

Case a will be considered in this section. Case b for $\mathfrak{g} = \mathfrak{sl}_2$ is considered in [5]. The case of a classical Lie algebra will be treated by our means elsewhere. The algebra $W = \mathbb{C}\langle x, y\rangle[[\hbar]]/\langle yx - xy - \hbar\rangle$ is called the *Weyl algebra*, it is isomorphic to the canonical deformation quantization of $(\mathbb{R}^2, \mathrm{d}x\wedge\mathrm{d}y)$ if we consider only polynomial functions on $\mathbb{R}^2$. This algebra admits a natural representation as indicated in the

Proposition 29. *The map* $\rho : W \longrightarrow \mathrm{End}_{\mathbb{C}[[\hbar]]}(\mathbb{C}[x][[\hbar]])$ *given by* $\rho(x)(f) = xf$ *and* $\rho(y)(f) = \hbar\frac{\partial f}{\partial x}$*, for any* $f \in \mathbb{C}[x][[\hbar]]$ *defines an irreducible representation of the Weyl algebra.*

We order the letters of the Weyl algebra as follows: $x < y < \hbar$. Assume we are given $A_i = (a_i, b_i) \in \mathbb{N}^2$, for $i \in [n]$. Set $A = (A_1, \dots, A_n) \in (\mathbb{N}^2)^n$, $X^{A_i} = x^{a_i}y^{b_i}$ and let $|\ | : \mathbb{N}^n \longrightarrow \mathbb{N}$ be the function such that $|x| := \sum_{i=1}^n x_i$, for all $x \in \mathbb{N}^n$. Given $x \in \mathbb{N}^n$ and $i \in \mathbb{N}$, we denote by $x_{<i}$ the vector $(x_1, \dots, x_{i-1}) \in \mathbb{N}^{i-1}$, by $x_{\leq i}$ the vector $(x_1, \dots, x_i) \in \mathbb{N}^i$ and by $x_{>i}$ the vector $(x_{i+1}, \dots, x_n) \in \mathbb{N}^{n-i}$. We write $a \vdash n$ if $a \in \mathbb{N}^k$ for some k and $|a| = n$. Using this notation we have

Definition 30. *The* normal coordinates $N(A,k)$ *of* $\prod_{i=1}^{n} X^{A_i} \in W$ *are defined through the identity*

$$\prod_{i=1}^{n} X^{A_i} = \sum_{k=0}^{\min} N(A,k) x^{|a|-k} y^{|b|-k} \hbar^k \tag{6}$$

for $0 \leq k \leq \min = \min(|a|,|b|)$*. For* $k > \min$*, we set* $N(A,k)$ *equal to* 0*.*

Recall that given finite sets N and M with n and m elements respectively, the number of one-to-one functions $f : N \longrightarrow M$ is $m(m-1)\ldots(m-n+1) = (m)_n$. The number $(m)_n$ is called the n-th *falling factorial* of m. The n-th *rising factorial* $m^{(n)}$ of m is given by $m^{(n)} = m(m+1)\ldots(m+n-1)$. For any $a, b \in \mathbb{N}^n$, we define $\binom{a}{b} := \binom{a_1}{b_1}\binom{a_2}{b_2}\ldots\binom{a_n}{b_n}$, and $a! = a_1!a_2!\ldots a_n!$.

Definition 31. *A k-pairing from set E to set F is an injective function from a k-elements subset of E to F. We denote by* $P_k(E,F)$ *the set of k-pairings from E to F.*

Definition 32. *Fix variables* $t = (t_1,\ldots,t_n)$ *and* $s = (s_1,\ldots,s_n)$*. The* generating series N *of the normal coordinates in the Weyl algebra is given by*

$$N = \sum_{a,b,c} N(A,c) \frac{s^a}{a!}\frac{t^b}{b!} u^c \in \mathbb{C}[[s,t,u]] \tag{7}$$

where the sum runs over $a, b \in \mathbb{N}^n$*,* $c \in \mathbb{N}$ *and* $A = (A_1,\ldots,A_n) \in (\mathbb{N}^2)^n$*.*

Theorem 33. *Let* A, k *be as in the Definition 30, the following identity holds*

a. $N(A,k) = \sum_{p \vdash k} \binom{b}{p} \prod_{i=1}^{n-1} (|a_{>i}| - |p_{>i}|)_{p_i}$*, where* $p \in \mathbb{N}^{n-1}$*.*

b. Let $E_1,\ldots,E_n,F_1,\ldots,F_n$ *be disjoint sets such that* $\sharp(E_i) = a_i$*,* $\sharp(F_i) = b_i$*, for* $i \in [n]$*, set* $E = \cup_{i=1}^{n} E_i$*, and* $F = \cup_{i=1}^{n} F_i$*, then* $N(A,k) = \sharp(\{p \in P_k(E,F) \mid \text{if } (a,p(a)) \in E_i \times F_j \text{ then } i > j\})$*.*

c. $N = \exp\left(\sum_{i>j} u t_i s_j + \sum_i t_i + \sum_j s_j\right)$*.*

Proof. Using induction one show that the following identity hold in the Weyl algebra

$$y^b x^a = \sum_{k=0}^{\min} \binom{b}{k} (a)_k x^{a-k} y^{b-k} \hbar^k, \tag{8}$$

where $\min = \min(a,b)$. Several applications of identity (8) imply a. Notice that for given sets E, F such that $\sharp(E) = a$ and $\sharp(F) = b$, $\binom{b}{k}(a)_k$ is equal to $\sharp(\{p \in P_k(E,F)\})$, showing part b for $n = 2$. The general formula follows from induction. This prove b. It follows from standard combinatorial facts (see [20], [22] for more details) that

$$\exp\left(\sum_{i>j} u t_i s_j + \sum_i t_i + \sum_j s_j\right) = \sum_{a,b,c} c_{a,b,c} \frac{s^a}{a!}\frac{t^b}{b!}\frac{u^c}{c!}$$

where $c_{a,b,c} = \sharp(\{(p,\sigma) : p \in P_c(A,B) \text{ and } \sigma : [c] \to p, \text{ a bijection }\})$ which is equivalent to formula (7). □

Figure 4 illustrates the combinatorial interpretation of the normal coordinate $N(A,k)$ of an element $\prod_{i=1}^{3} X^{A_i} \in W$, it shows are of the possible pairing contributing to $N(A,k)$.



Figure 4: Combinatorial interpretation of N.

Corollary 34. *For any given $(t,a,b) \in \mathbb{N} \times \mathbb{N}^n \times \mathbb{N}^n$, the following identity holds*

$$\prod_{i=1}^{n} (t + |a_{>i}| - |b_{>i}|)_{b_i} = \sum_{p \vdash k} \binom{b}{p} \prod_{i=1}^{n-1} (|a_{>i}| - |p_{>i}|)_{p_i} \, t_{|b|-k}$$

Proof. Consider the identity (6) in the representation of the Weyl algebra defined in Proposition 29. Apply both sides of the identity (6) to x^t for $t \in \mathbb{N}$ and use Theorem 33. ☐

Theorem 35. *Let A, c be as in the Definition 30. The following identity holds*

$$N(A,c) = \sum_{\sum c_{ij} = c} \prod_{j=2}^{k} \binom{a_j}{\sum_i c_{ij}} \binom{\sum_i c_{ij}}{c_{ij} \dots c_{(j-1)j}} \prod_{i=1}^{k-1} \binom{b_i}{\sum_j c_{ij}} \binom{\sum_j c_{ij}}{c_{i(i+1)} \dots c_{ik}} \prod_{i>j} c_{ij}!$$

where c_{ij} are integers with $1 \leq i \leq k-1$, $2 \leq j \leq k$ and $i > j$.

Proof. Consider maps $F_{ij} : \{p \in P_c(E,F) \mid \text{if } (a, p(a)) \in E_i \times F_j \text{ then } i > j\} \longrightarrow \mathbb{N}$ given by $F_{ij}(p) = \sharp(\{(a,p(a)) \in E_i \times F_j\})$. Notice that $N(A,c) = \sum_{\sum c_{ij} = c} \sharp(\{p \in P_c(E,F) : F_{ij}(p) = c_{ij}, i > j\})$. Moreover

$$\sharp(\{p \in P_c(E,F) : F_{ij}(p) = c_{ij}, i > j\}) = \prod_{j=2}^{k} \binom{a_j}{\sum_i c_{ij}} \binom{\sum_i c_{ij}}{c_{ij} \dots c_{(j-1)j}} \prod_{i=1}^{k-1} \binom{b_i}{\sum_j c_{ij}} \binom{\sum_j c_{ij}}{c_{i(i+1)} \dots c_{ik}} \prod_{i>j} c_{ij}!.$$

☐

6.2 Quantum symmetric functions of Weyl type

The following theorem provides explicit formula for the product of m elements of the n-th symmetric power of the Weyl algebra. Let us explain our notation: fix a matrix $A : [m] \times [n] \longrightarrow \mathbb{N}^2$, $(A_{ij}) = ((a_{ij}), (b_{ij}))$. Given $\sigma \in (S_n)^m$ and $j \in [n]$, A^{σ}_j denotes the vector $(A_{1\sigma_1^{-1}(j)}, \dots, A_{m\sigma_m^{-1}(j)}) \in (\mathbb{N}^2)^n$ and set $X_j^{A_{ij}} = x_j^{a_{ij}} y_j^{b_{ij}}$ for $j \in [n]$. Set $|A^{\sigma}_j| = (|a^{\sigma}_j|, |b^{\sigma}_j|)$ where $|a^{\sigma}_j| = \sum_{i=1}^{m} a_{i\sigma_i^{-1}(j)}$ and $|b^{\sigma}_j| = \sum_{i=1}^{m} b_{i\sigma_i^{-1}(j)}$. We have the following

Theorem 36. *For any $A : [m] \times [n] \longrightarrow \mathbb{N}^2$, the following identity*

$$(n!)^{m-1} \prod_{i=1}^{m} \left(\overline{\prod_{j=1}^{n} X_j^{A_{ij}}} \right) = \sum_{\sigma,k,p} \left(\prod_{i,j} \binom{b^{\sigma}_j}{p^j} (|(a^{\sigma}_j)_{>i}| - |p^j_{>i}|)_{p^j_i} \right) \overline{\prod_{j=1}^{n} X_j^{|A^{\sigma}_j| - (k_j, k_j)}} \hbar^{|k|} \tag{9}$$

where $\sigma \in \{\mathrm{id}\} \times S_n^{m-1}$, $k \in \mathbb{N}^n$, $(i,j) \in [m-1] \times [n]$ and $p = p^j_i \in (\mathbb{N}^{m-1})^n$, holds in $\mathrm{Sym}^n(W)$.

Proof. We use Theorem 14 and Theorem 33

$$\begin{aligned}
(n!)^{m-1}\overline{\prod_{i=1}^{m}\left(\prod_{j=1}^{n} X_j^{A_{ij}}\right)} &= \sum_{\sigma\in\{\mathrm{id}\}\times S_n^{m-1}} \overline{\prod_{j=1}^{n}\left(\prod_{i=1}^{m} X_j^{A_{i\sigma_i^{-1}(j)}}\right)}\\
&= \sum_{\sigma\in\{\mathrm{id}\}\times S_n^{m-1}} \overline{\prod_{j=1}^{n}\left(\sum_{k=0}^{\min_j} N(A_j^\sigma,k) X_j^{|A_j^\sigma|-(k,k)}\hbar^k\right)}\\
&= \sum_{\sigma,k}\left(\prod_{j=1}^{n} N(A_j^\sigma,k_j)\right)\overline{\prod_{j=1}^{n} X_j^{|A_j^\sigma|-(k_j,k_j)}}\hbar^{|k|}\\
&= \sum_{\sigma,k,p}\left(\prod_{i,j}\binom{b_j^\sigma}{p^j}(|(a_j^\sigma)_{>i}|-|p_{>i}^j|)_{p_i^j}\right)\overline{\prod_{j=1}^{n} X_j^{|A_j^\sigma|-(k_j,k_j)}}\hbar^{|k|},
\end{aligned}$$

where $\min_j = \min(|a_j^\sigma|,|b_j^\sigma|)$. ☐

Now we proceed to state and prove the quantum analogue of Theorem 22. We begin by introducing a $\star$-product on $(\mathbb{C}[\mathbb{N}^{2m}]^{\otimes n}[[\hbar]])_{S_n}$, which is motivated by the proof of Theorem 39 below.

Definition 37. *The $\star$-product on $(\mathbb{C}[\mathbb{N}^{2m}]^{\otimes n}[[\hbar]])_{S_n}$ for $A\in(\mathbb{N}^{2m})^n$ and $C\in(\mathbb{N}^{2m})^n$ is given by the formula*

$$\overline{A}\star\overline{C} = \frac{1}{n!}\sum_{I,\sigma}\binom{b}{I}(\sigma(c))_I\overline{A+\sigma(C)-(I,I)}\hbar^{|I|} \tag{10}$$

where $I:[n]\times[m]\to\mathbb{N}$ and $\sigma\in S_n$.

Definition 38. *Fix $m\in\mathbb{N}$. The algebra of quantum symmetric functions of type A_n, B_n and D_n are given by*

- $\mathrm{QSym}_{A_n}(m) = (\mathbb{C}[x_1,y_1,\ldots,x_n,y_n][[\hbar]],\star)_{S_n} \cong (\mathbb{C}[x_1,y_1,\ldots,x_n,y_n][[\hbar]],\star)^{S_n}$,
- $\mathrm{QSym}_{B_n}(m) = (\mathbb{C}[x_1,y_1,\ldots,x_n,y_n][[\hbar]],\star)_{\mathbb{Z}_2^n\rtimes S_n} \cong (\mathbb{C}[x_1,y_1,\ldots,x_n,y_n][[\hbar]],\star)^{\mathbb{Z}_2^n\rtimes S_n}$,
- $\mathrm{QSym}_{D_n}(m) = (\mathbb{C}[x_1,y_1,\ldots,x_n,y_n][[\hbar]],\star)_{\mathbb{Z}_2^{n-1}\rtimes S_n} \cong (\mathbb{C}[x_1,y_1,\ldots,x_n,y_n][[\hbar]],\star)^{\mathbb{Z}_2^{n-1}\rtimes S_n}$,

where $x_i=(x_{i1},\ldots,x_{im})$ and $y_i=(y_{i1},\ldots,y_{im})$.

Theorem 39 (Quantum symmetric functions).

a. The following is a commutative diagram

$$\begin{tikzcd}
(\mathbb{C}[\mathbb{N}^{2m}]^{\otimes n}[[\hbar]],\star)_{S_n} \arrow[r, phantom, "\supset"] \arrow[d] & (\mathbb{C}[\mathbb{N}_e^{2m}+\mathbb{N}_o^{2m}]^{\otimes n}[[\hbar]],\star)_{S_n} \arrow[r, phantom, "\supset"] \arrow[d] & (\mathbb{C}[\mathbb{N}_e^{2m}]^{\otimes n}[[\hbar]],\star)_{S_n} \arrow[d] \\
\mathrm{QSym}_{A_n}(m) \arrow[r, two heads] & \mathrm{QSym}_{D_n}(m) \arrow[r, two heads] & \mathrm{QSym}_{B_n}(m)
\end{tikzcd}$$

where the vertical arrows are isomorphisms.

b. For $A_i = (a_i, b_i) \in (\mathbb{N}^2)^m$ and $C_i = (c_i, d_i) \in (\mathbb{N}^2)^m$ $i \in [n]$, we set $X_i^{A_i} = x_i^{a_i} y_i^{b_i}$ and $X_i^{C_i} = x_i^{c_i} y_i^{d_i}$ where $x_i = (x_{i1}, \ldots, x_{im})$ and $y = (y_{i1}, \ldots, y_{im})$, the product rule in $\mathrm{QSym}_{A_n}(m)$ is given by

$$\overline{X_1^{A_1} X_2^{A_2} \ldots X_n^{A_n}} \star \overline{X_1^{C_1} X_2^{C_2} \ldots X_n^{C_n}} = \frac{1}{n!} \sum_{I,\sigma} \binom{b}{I} (\sigma(c))_I \overline{X^{A+\sigma(C)-(I,I)}} \hbar^{|I|},$$

where $I : [n] \times [m] \to \mathbb{N}$, $\sigma \in S_n$. The product on $\mathrm{QSym}_{B_n}(m)$ and $\mathrm{QSym}_{D_n}(m)$ is given by restriction.

Proof. We prove b which implies a. Given $A_i, C_i \in (\mathbb{N}^2)^m$, using (10), we obtain

$$\begin{aligned}
\overline{X_1^{A_1} X_2^{A_2} \ldots X_n^{A_n}} \star \overline{X_1^{C_1} X_2^{C_2} \ldots X_n^{C_n}} &= \frac{1}{n!} \sum_{\sigma \in S_n} \overline{X_1^{A_1} X_1^{C_{\sigma^{-1}(1)}} X_2^{A_2} X_2^{C_{\sigma^{-1}(2)}} \ldots X_n^{A_n} X_n^{C_{\sigma^{-1}(n)}}} \\
&= \frac{1}{n!} \sum_{\sigma \in S_n} \overline{x_1^{a_1} y_1^{b_1} x_1^{c_{\sigma^{-1}(1)}} y_1^{d_{\sigma^{-1}(1)}} \ldots x_n^{a_n} y_n^{b_n} x_n^{c_{\sigma^{-1}(n)}} y_n^{d_{\sigma^{-1}(n)}}} \\
&= \frac{1}{n!} \sum_{\sigma \in S_n} \left(\prod_{j=1}^{n} \sum_{i_j=0}^{\min_j} \binom{b_j}{i_j} (c_{\sigma^{-1}(j)})_{i_j} \overline{X_j^{A_j + C_{\sigma^{-1}(j)} - (i_j, i_j)}} \hbar^{i_j} \right) \\
&= \frac{1}{n!} \sum_{I,\sigma} \left(\binom{b}{I} (\sigma(c))_I \right) \overline{X^{A+\sigma(C)-(I,I)}} \hbar^{|I|}
\end{aligned}$$

where $\min_j = \min(b_j, c_{\sigma^{-1}(j)})$. Now, consider $A_i, C_i \in \mathbb{N}_e^{2m}$

$$\begin{aligned}
\overline{X_1^{A_1} \ldots X_n^{A_n}} \star \overline{X_1^{C_1} \ldots X_n^{C_n}} &= \frac{1}{2^n n!} \sum_{(t,\sigma) \in \mathbb{Z}_2^n \rtimes S_n} \overline{(x_1^{a_1} y_1^{b_1} \ldots x_n^{a_n} y_n^{b_n})((t,\sigma) x_1^{c_1} y_1^{d_1} \ldots x_n^{c_n} y_n^{d_n})} \\
&= \frac{1}{2^n n!} \sum_{I,\sigma} \left(\sum_{t \in \mathbb{Z}_2^n} (-1)^{\sum_{t_i=-1} c_i + d_i} \binom{b}{I} (\sigma(c))_I \right) \overline{X^{A+\sigma(C)-(I,I)}} \hbar^{|I|} \\
&= \frac{1}{n!} \sum_{I,\sigma} \left(\binom{b}{I} (\sigma(c))_I \right) \overline{X^{A+\sigma(C)-(I,I)}} \hbar^{|I|}
\end{aligned}$$

since $\sum_{t \in \mathbb{Z}_2^n} (-1)^{\sum_{t_i=-1} c_i + d_i} = 2^n$. Finally, consider $A_i, C_i \in \mathbb{N}_e^{2m} + \mathbb{N}_o^{2m}$

$$\begin{aligned}
\overline{X_1^{A_1} \ldots X_n^{A_n}} \star \overline{X_1^{C_1} \ldots X_n^{C_n}} &= \frac{1}{2^{n-1} n!} \sum_{(t,\sigma) \in \mathbb{Z}_2^n \rtimes S_n} \overline{(x_1^{a_1} y_1^{b_1} \ldots x_n^{a_n} y_n^{b_n})(t,\sigma) x_1^{c_1} y_1^{d_1} \ldots x_n^{c_n} y_n^{d_n}} \\
&= \frac{1}{2^{n-1} n!} \sum_{I,\sigma} \left(k(t,c,d) \binom{b}{I} (\sigma(c))_I \right) \overline{X^{A+\sigma(C)-(I,I)}} \hbar^{|I|} \\
&= \frac{1}{n!} \sum_{I,\sigma} \left(\binom{b}{I} (\sigma(c))_I \right) \overline{X^{A+\sigma(C)-(I,I)}} \hbar^{|I|}
\end{aligned}$$

since $k(t,c,d) = \sum_{\substack{t\in\mathbb{Z}_2^n \\ \prod t_i=1}} (-1)^{\sum_{t_i=-1} c_i+d_i+c_n+d_n} = 2^{n-1}$ ☐

6.3 Quantum symmetric functions on $\mathbb{C}^n/\mathbb{Z}_m^n \rtimes S_n$

In this section we shift to complex analytic notation to study the Poisson orbifolds $\mathbb{C}^n/\mathbb{Z}_m^n \rtimes S_n$ and $\mathbb{C}^n/\mathcal{D}_m^n \rtimes S_n$. The deformation quantization of $\mathbb{C}^n$ provided with the canonical symplectic structure is isomorphic to the Weyl algebra $W = \mathbb{C}\langle z_1, \overline{z_1}, \dots, z_n, \overline{z_n}\rangle[[\hbar]]/\langle z\overline{z} - \overline{z}z - 2i\hbar\rangle$. The group $\mathbb{Z}_m^n \rtimes S_n$ acts on $\mathbb{C}^n$ as follows

$$\begin{array}{ccc} (\mathbb{Z}_m^n \rtimes S_n) \times \mathbb{C}^n & \longrightarrow & \mathbb{C}^n \\ ((w_1,\dots,w_n),\sigma)(z_1,\dots,z_n) & \longmapsto & (w_1 z_{\sigma^{-1}(1)},\dots,w_n z_{\sigma^{-1}(n)}) \end{array}$$

where $w_j = e^{\frac{2\pi i k_j}{m}}$, $j = 1,\dots,n$. Thus $\mathbb{Z}_m^n \rtimes S_n$ acts on $W = \mathbb{C}\langle z_1, \overline{z_1}, \dots, z_n, \overline{z_n}\rangle[[\hbar]]/\langle z\overline{z} - \overline{z}z - 2i\hbar\rangle$. We denote $Z_i^{A_i} = z_i^{a_i}\overline{z}_i^{b_i}$ and $\mathbb{N}_m^2 = \{(a,b) :$ there is $k \in \mathbb{Z}$ such that $b - a = km\}$.

Definition 40. *The $\star$-product on $(\mathbb{C}[\mathbb{N}_m^2]^{\otimes n}[[\hbar]])_{\mathbb{Z}_m^n \rtimes S_n}$ for $A \in (\mathbb{N}^2)^n$ and $C \in (\mathbb{N}^2)^n$ is given by the formula*

$$\overline{A} \star \overline{C} = \frac{1}{n!}\sum_{I,\sigma} (-2i)^{|I|}\binom{b}{I}(\sigma(c))_I \overline{A + \sigma(C) - (I,I)}\hbar^{|I|}$$

where $I : [n] \to \mathbb{N}$ and $\sigma \in S_n$.

Theorem 41. *The map*

$$\begin{array}{ccc} (\mathbb{C}[\mathbb{N}_m^2]^{\otimes n}[[\hbar]], \star)_{\mathbb{Z}_m^n \rtimes S_n} & \longrightarrow & (\mathbb{C}[z,\overline{z}]^{\otimes n}[[\hbar]], \star)_{\mathbb{Z}_m^n \rtimes S_n} \\ (A_1,\dots,A_n) & \longmapsto & Z_1^{A_1}\dots Z_n^{A_n} \end{array}$$

is an algebra isomorphism.

Proof. Let $A_i = (a_i, b_i) \in \mathbb{N}^2$, and $C_i = (c_i, d_i) \in \mathbb{N}^2$ $i \in [n]$, we have

$$\begin{aligned} \overline{Z_1^{A_1}\dots Z_n^{A_n}} \star \overline{Z_1^{C_1}\dots Z_n^{C_n}} &= \overline{z_1^{a_1}\overline{z}_1^{b_1}\dots z_n^{a_n}\overline{z}_n^{b_n}}\ \overline{z_1^{c_1}\overline{z}_1^{d_1}\dots z_n^{c_n}\overline{z}_n^{d_n}} \\ &= \frac{1}{m^n n!}\sum_{(w,\sigma)\in\mathbb{Z}_m^n\rtimes S_n} \overline{z_1^{a_1}\overline{z}_1^{b_1}\dots z_n^{a_n}\overline{z}_n^{b_n}((w,\sigma)(z_1^{c_1}\overline{z}_1^{d_1}\dots z_n^{c_n}\overline{z}_n^{d_n}))} \\ &= \frac{1}{m^n n!}\sum_{\sigma\in S_n}\left(\prod_{j=1}^{n}\sum_{k_j=0}^{m-1}(e^{\frac{2\pi i}{m}(d_j-c_j)})^{k_j}\right)\overline{\prod_{j=1}^{n} z_j^{a_j}\overline{z}_j^{b_j} z_j^{c_{\sigma^{-1}(j)}}\overline{z}_j^{d_{\sigma^{-1}(j)}}} \\ &= \frac{1}{n!}\sum_{\substack{I:[n]\to\mathbb{N} \\ \sigma\in S_n}}\left((-2i)^{|I|}\binom{b}{I}(\sigma(c))_I\right)\overline{Z^{A+\sigma(C)-(I,I)}}\hbar^{|I|} \end{aligned}$$

☐

Let $\mathcal{D}_m$ be the dihedral group of order $2n$, where $\mathcal{D}_m = \{R_0, R_1, \ldots, R_{m-1}, S_0, S_1, \ldots, S_{m-1}\}$, $R_k(z) = e^{\frac{2\pi ik}{n}} z$, and $S_k(z) = e^{\frac{2\pi ik}{n}} \overline{z}$, for $k = 0, \ldots, m-1$. $\mathcal{D}_m^n \rtimes S_n$ acts on $\mathbb{C}^n$ as follows

$$\begin{array}{rcl} (\mathcal{D}_m^n \rtimes S_n) \times \mathbb{C}^n & \longrightarrow & \mathbb{C}^n \\ ((d_1, d_2, \ldots, d_n), \sigma)(z_1, z_2, \ldots, z_n) & \longmapsto & (d_1 z_{\sigma^{-1}(1)}, \ldots, d_n z_{\sigma^{-1}(n)}) \end{array}$$

We will apply Polya functor to obtain a product rule on $(C^\infty(\mathbb{C}^n)[[\hbar]], \star)_{\mathcal{D}^n \rtimes S_n}$

Definition 42. *The $\star$-product on $(\mathbb{C}[\mathbb{N}_m^2]^{\otimes n}[[\hbar]], \star)_{\mathcal{D}^n \rtimes S_n}$ for $A \in (\mathbb{N}^2)^n$ and $C \in (\mathbb{N}^2)^n$ is given by the formula*

$$\overline{A} \star \overline{C} = \frac{1}{2n!} \sum_{I,\sigma} \binom{b}{I} (\sigma(c))_I \overline{A + (\sigma(c), \sigma(d)) - (I, I)} \hbar^{|I|} + \frac{1}{2n!} \sum_{I,\sigma} \binom{b+c}{I} (\sigma(d))_I \overline{A + (\sigma(d), \sigma(c)) - (I, I)} \hbar^{|I|}$$

where $I : [n] \to \mathbb{N}$ and $\sigma \in S_n$.

Theorem 43. *The map $(\mathbb{C}[\mathbb{N}_m^2]^{\otimes n}[[\hbar]], \star)_{\mathcal{D}^n \rtimes S_n} \longrightarrow (\mathbb{C}[z, \overline{z}]^{\otimes n}[[\hbar]], \star)_{\mathcal{D}^n \rtimes S_n}$ given by $(A_1, \ldots, A_n) \longmapsto Z_1^{A_1} \ldots Z_n^{A_n}$ is an algebra isomorphism.*

Proof. For any $A_i = (a_i, b_i) \in \mathbb{N}^2$, $C_i = (c_i, d_i) \in \mathbb{N}^2$, we have

$$\begin{array}{l} \overline{Z_1^{A_1} \ldots Z_n^{A_n}}\ \overline{Z_1^{C_1} \ldots Z_n^{C_n}} \\ = \quad \displaystyle\frac{1}{(2n)n!} \sum_{(d,\sigma) \in \mathcal{D}^n \rtimes S_n} \overline{z_1^{a_1} \overline{z}_1^{b_1} \ldots z_n^{a_n} \overline{z}_n^{b_n} ((d, \sigma) z_1^{c_1} \overline{z}_1^{d_1} \ldots z_n^{c_n} \overline{z}_n^{d_n})} \\ = \quad \displaystyle\frac{1}{(2n)n!} \sum_{\sigma \in S_n} \left(\prod_{j=1}^n \sum_{k_j=0}^{n-1} e^{\frac{2\pi i k_j}{n}(d_j - c_j)} \right) \left(\prod_{j=1}^n \overline{z_j^{a_j} \overline{z_j}^{b_j} z_j^{c_{\sigma^{-1}(j)}} \overline{z_j}^{d_{\sigma^{-1}(j)}}} + \overline{z_j^{a_j} \overline{z_j}^{b_j + c_{\sigma^{-1}(j)}} z_j^{d_{\sigma^{-1}(j)}}} \right) \\ = \quad \displaystyle\frac{1}{2n!} \sum_{\sigma,I} \binom{b}{I} (\sigma(c))_I \overline{Z^{A + (\sigma(c), \sigma(d)) - (I,I)}} \hbar^{|I|} + \frac{1}{2n!} \sum_{\sigma,I} \binom{b+c}{I} (\sigma(d))_I \overline{Z^{A + (\sigma(d), \sigma(c)) - (I,I)}} \hbar^{|I|}. \end{array}$$

where $\sigma \in S_n$ and $I : [n] \to \mathbb{N}$. □

6.4 Quantum super-functions

We denote by $\mathrm{Mat}(n)$ the algebra of $\mathbb{C}$-matrices of order $n \times n$. It is well-known see [3] that the canonical quantization of $\bigwedge_{\mathbb{C}}[\theta_1, \ldots, \theta_m]$ is isomorphic to the Clifford algebra $C(m)$, i.e, the complex free algebra on m generators $\theta_1, \ldots, \theta_m$ subject to the relations $\theta_i \theta_j + \theta_j \theta_i = 2\delta_{ij}$, for $i, j \in [m]$. It is also known that

$$\begin{array}{rl} C(m) \cong & \mathrm{Mat}(2^{\frac{m}{2}}) \text{ if } m \text{ is even, and} \\ & \\ C(m) \cong & \mathrm{Mat}(2^{\frac{m-1}{2}}) \oplus \mathrm{Mat}(2^{\frac{m-1}{2}}) \text{ if } m \text{ is odd.} \end{array}$$

Thus the algebra of quantum symmetric functions on the super-space $\mathbb{R}^{0|m}$ is isomorphic to

$$\begin{array}{rcl} \mathrm{Sym}^n(\mathrm{Mat}(2^m)) & \cong & \mathrm{Schur}(n, 2^m) \text{ for } m \text{ even,} \\ \mathrm{Sym}^n(\mathrm{Mat}(2^{\frac{m-1}{2}}) \oplus \mathrm{Mat}(2^{\frac{m-1}{2}})) & \cong & \displaystyle\bigoplus_{a+b=n} \mathrm{Sym}^a(\mathrm{Mat}(2^{\frac{m-1}{2}})) \otimes \mathrm{Sym}^b(\mathrm{Mat}(2^{\frac{m-1}{2}})) \\ & \cong & \displaystyle\bigoplus_{a+b=n} \mathrm{Schur}(a, 2^{\frac{m-1}{2}}) \otimes \mathrm{Schur}(b, 2^{\frac{m-1}{2}}) \text{ for } m \text{ odd.} \end{array}$$

Definition 44 and Proposition 45 below are taken from [18].

Definition 44. *We denote by* $\mathfrak{gl}(\infty)$ *the algebra of all matrices* (a_{ij}) *such that* $a_{ij} \in \mathbb{C}[[\hbar]]$ *if* $i \geq j$, *and* $a_{ij} \in (\hbar\mathbb{C}[[\hbar]])^{j-i}$, *if* $i < j$. *We set* $\mathrm{Schur}(\infty, n) := \mathrm{Sym}^n(\mathfrak{gl}(\infty))$.

For any $a, b \in \mathbb{Z}^{\geq 0}$, we define the matrices

$$E_{a,b}(i,j) = \begin{cases} \frac{(b+k)!}{k!}\hbar^b, & \text{if } (i,j) = (a,b) + (k,k),\ k = 0,1,2,\ldots \\ \\ 0, & \text{otherwise.} \end{cases}$$

Proposition 45. *There is a canonical isomorphism* $\rho : W \longrightarrow \mathfrak{gl}(\infty)$ *defined on generators by* $\rho(x) = E_{1,0}$, $\rho(y) = E_{0,1}$ *and* $\rho(\hbar) = \hbar I$.

Proof. The linear map ρ is a well defined algebra homomorphism since the Weyl algebra is the quotient of a (formal) free algebra by the ideal generated by the relation $yx = xy + \hbar$, and the following identity holds $E_{0,1}E_{1,0} = E_{1,0}E_{0,1} + \hbar I$ in $\mathfrak{gl}(\infty)$. Similarly ρ is a bijection since $E_{a,b}$ is a basis for $\mathfrak{gl}(\infty)$ and $\rho(x^a y^b) = E_{a,b}$ as consequence of the fact that in $\mathfrak{gl}(\infty)$ the following identities are satisfied: $E_{0,a}E_{0,1} = E_{0,a+1}$, $E_{a,0}E_{1,0} = E_{a+1,0}$ and $E_{a,0}E_{0,b} = E_{a,b}$, for all $a, b \in \mathbb{Z}^{\geq 0}$.

□

We denote by $\widehat{W}$ the formal Weyl algebra, i.e. $\widehat{W} = \mathbb{C}\langle\langle x, y\rangle\rangle[[\hbar]]/\langle yx - xy - \hbar\rangle$ and $\hat{\mathrm{Q}}\mathrm{Sym}_{A_n}(1) = \mathrm{Sym}^n(\widehat{W})$.

Theorem 46. a. *The algebra* $\hat{\mathrm{Q}}\mathrm{Sym}_{A_n}(1)$ *of formal quantum symmetric functions on* $((\mathbb{R}^2)^n, \sum \mathrm{d}x_i \wedge \mathrm{d}y_i)$ *is isomorphic to* $\mathrm{Schur}(\infty, n)$.

b. *The algebra of formal quantum symmetric functions on the superspaces* $\mathbb{R}^{2|n}$ *is isomorphic to* $\mathrm{Sym}^n(\mathfrak{gl}(\infty) \otimes C(m))$.

Proof. a. $\hat{\mathrm{Q}}\mathrm{Sym}_{A_n}(1) \cong \mathrm{Sym}^n(\widehat{W}) \cong \mathrm{Schur}(\infty, n)$.

b. The algebra of formal quantum functions on the superspace $\mathbb{R}^{2|n}$ is isomorphic to

$$\widehat{W} \otimes C(m) \cong \mathfrak{gl}(\infty) \otimes C(m).$$

Thus, the algebra of formal quantum symmetric function on the superspaces $\mathbb{R}^{2m|n}$ is isomorphic to $\mathrm{Sym}^n(\mathfrak{gl}(\infty) \otimes C(m))$.

□

6.5 M-Weyl algebra

In this section we introduce the M-Weyl algebra. Although closely related to the Weyl algebra, the M-Weyl algebra does not arises as an instance of the Kontsevich star product.

Definition 47. *The* M-Weyl algebra *is the algebra* $MW = \mathbb{C}\langle x, y\rangle[[\hbar]]/\langle yx - xy - x^2\hbar\rangle$. *The letter* M *stands for meromorphic or mimetic.*

We have the following analogue of Proposition 29

Proposition 48. *The map* $\rho : MW \longrightarrow \mathrm{End}(\mathbb{C}[x][[\hbar]])$ *given by* $\rho(x)(f) = x^{-1}f$ *and* $\rho(y)(f) = -\hbar\frac{\partial f}{\partial x}$, *for any* $f \in \mathbb{C}[x][[\hbar]]$ *defines a representation of the* M*-Weyl algebra.*

We order the letters of the M-Weyl algebra as follows: $x < y < \hbar$. Assume we are given $A_i = (a_i, b_i) \in \mathbb{N}^2$, for $i \in [n]$. Set $A = (A_1, \ldots, A_n) \in (\mathbb{N}^2)^n$ and $X^{A_i} = x^{a_i}y^{b_i}$, for $i \in [n]$. Using this notation we have

Definition 49. *The* normal coordinates $N_M(A, k)$ *of* $\prod_{i=1}^{n} X^{A_i} \in MW$ *are defined through the identity*

$$\prod_{i=1}^{n} X^{A_i} = \sum_{k=0}^{\min} N_M(A, k)x^{|A|+(k,-k)}\hbar^k \tag{11}$$

where $0 \leq k \leq \min = \min(|a|, |b|)$. *For* $k > \min$, *we set* $N_M(A, k) = 0$.

Theorem 50. *Let* A, k *be as in the previous definition, the following identity holds*

a.

$$N_M(A, k) = \sum_{p \vdash k} \binom{b}{p} \prod_{i=1}^{n-1} (|a_{>i}| + |p_{>i}|)^{(p_i)}. \tag{12}$$

b. Let $E_1, \ldots, E_n, F_1, \ldots, F_n$ *be disjoint sets such that* $\sharp(F_i) = a_i$, $\sharp(E_i) = b_i$, *for* $i \in [n]$. *Set* $E = \cup E_i$, $F = \cup F_i$ *and consider the set* M_k *of all functions* $f : F \longrightarrow P(E)$ *such that*

- $f(x) \cap f(y) = \emptyset$, *for all* $x, y \in F$.
- *If* $x \in F_i$, $y \in E_j$, *and* $y \in f(x)$, *then* $j < i$.
- $\sum_{a \in F} \sharp(f(a)) = k$.

then $N_M(A, k) = \sharp(M_k)$.

Proof. Using induction one show that the following identity hold in the M-Weyl algebra

$$y^b x^a = \sum_{k=0}^{\min} \binom{b}{k} a^{(k)} x^{a+k} y^{b-k} \hbar^k, \tag{13}$$

where $\min = \min(a, b)$. Several application of the identity (13) imply a. Notice that for given sets E, F such that $|F| = a$ and $|E| = b$, $\binom{b}{k}a^{(k)}$ is equal to $\sharp(\{f : F \longrightarrow P(E) : \ f(x) \cap f(y) = \emptyset,$ for all $x, y \in F,$ and $\sum_{a \in F} \sharp(f(a)) = k\})$. This shows b, for $n = 1$. The general formula follows from induction. $\square$

Figure 5 illustrate the combinatorial interpretation of the normal coordinates $N_M(A, 6)$ of $\prod_{i=1}^{3} X^{A_i} \in MW$, it shows an example of a function contributing to $N_M(A, 6)$.



Figure 5: Combinatorial interpretation of N_M.

Corollary 51. *For any given $(t, a, b) \in \mathbb{N} \times \mathbb{N}^n \times \mathbb{N}^n$, the following identity holds*

$$\prod_{i=1}^{n}(t - |a_{>i}| - |b_{>i}|)_{b_i} = \sum_{p \vdash k}(-1)^k \binom{b}{p} \prod_{i=1}^{n-1}(|a_{>i}| - |p_{>i}|)^{(p_i)} t_{|b|-k}$$

Proof. Consider the identity (11) in the representation of the M-Weyl algebra defined in Proposition 48. Apply both sides of the identity (11) to x^t and use Theorem 50 formula (12). $\square$

The following theorem provides explicit formula for the product of m elements of the M-Weyl algebra. Using the same notation as in the Theorem 36, we have the following

Theorem 52 (Symmetric powers of M-Weyl algebra). *For any $A : [m] \times [n] \longrightarrow \mathbb{N}^2$, the following identity*

$$(n!)^{m-1}\overline{\prod_{i=1}^{m}\left(\prod_{j=1}^{n} X_j^{A_{ij}}\right)} = \sum_{\sigma,k,p}\left(\prod_{i,j}\binom{b_j^\sigma}{p^j}(|(a_j^\sigma)_{>i}| + |p_{>i}^j|)^{(p_i^j)}\right)\overline{\prod_{j=1}^{n} X_j^{|A_j^\sigma| - (k_j,k_j)}}\hbar^{|k|} \tag{14}$$

where $\sigma \in \{\mathrm{id}\} \times S_n^{m-1}$, $k \in \mathbb{N}^n$, $(i,j) \in [m-1] \times [n]$ and $p = p_i^j \in (\mathbb{N}^{m-1})^n$, holds in $\mathrm{Sym}^n(MW)$.

Proof. By Theorem 14 and Theorem 50, we have

$$\begin{aligned}
(n!)^{m-1}\overline{\prod_{i=1}^{m}\left(\prod_{j=1}^{n} X_j^{A_{ij}}\right)} &= \sum_{\sigma \in \{\mathrm{id}\} \times S_n^{m-1}} \overline{\prod_{j=1}^{n}\left(\prod_{i=1}^{m} X_j^{A_{i\sigma_i^{-1}(j)}}\right)} \\
&= \sum_{\sigma \in \{\mathrm{id}\} \times S_n^{m-1}} \overline{\prod_{j=1}^{n}\left(\sum_{k=0}^{\min_j} N_M(A_j^\sigma, k) X_j^{|A_j^\sigma| + (k,-k)}\hbar^k\right)} \\
&= \sum_{\sigma,k}\left(\prod_{j=1}^{n} N_M(A_j^\sigma, k_j)\right)\overline{\prod_{j=1}^{n} X_j^{|A_j^\sigma| + (k_j,-k_j)}}\hbar^{|k|} \\
&= \sum_{\sigma,k,p}\left(\prod_{i,j}\binom{b_j^\sigma}{p^j}(|(a_j^\sigma)_{>i}| + |p_{>i}^j|)^{(p_i^j)}\right)\overline{\prod_{j=1}^{n} X_j^{|A_j^\sigma| - (k_j,k_j)}}\hbar^{|k|},
\end{aligned}$$

where $\min_j = \min(|a_j^\sigma|, |b_j^\sigma|)$. $\square$

6.6 Cohomological interpretation of symmetric functions

Let G be a finite group acting on a compact differentiable manifold X. G acts on $\mathrm{H}_\bullet(X)$, the singular homology groups with complex coefficients of X, as follows:

$$\begin{array}{ccc} G \times \mathrm{H}_\bullet(X) & \longrightarrow & \mathrm{H}_\bullet(X) \\ (g, \alpha) & \longmapsto & g_*(\alpha) \end{array}$$

where $g_*(\alpha)$ denotes the push-forward of α by the map g. Similarly G acts on $\mathrm{H}^\bullet(X)$, the singular cohomology groups with complex coefficients of X, as follows:

$$\begin{array}{ccc} G \times \mathrm{H}^\bullet(X) & \longrightarrow & \mathrm{H}^\bullet(X) \\ (g, \alpha) & \longmapsto & g^*(\alpha) \end{array}$$

where $g^*(\alpha)$ denotes the pull-back of α by the map g. It is well-known that $\mathrm{H}^\bullet(X/G) = \mathrm{H}^\bullet(X)^G$, see [11]. Identifying $\mathrm{H}^\bullet(X)^G$ with $\mathrm{H}^\bullet(X)_G$, we obtain $\mathrm{H}^\bullet(X/G) = \mathrm{H}^\bullet(X)_G$. Consider $X^n/G^n \rtimes K$ where $K \subset S_n$ and $X^n = X \times X \times \cdots \times X$. We have that

$$\mathrm{H}^\bullet(X^n/G^n \rtimes K) \cong \mathrm{H}^\bullet(X)^{\otimes n}/G^n \rtimes K \cong P_{G,K}(\mathrm{H}^\bullet(X)).$$

Next theorem shows how symmetric and supersymmetric functions arise as the cohomology groups of global orbifolds (quotient of manifolds by finite group actions). We denote by $\mathbb{CP}^\infty$ the inductive limit of the complex projective spaces $\mathbb{CP}^n$. Also, we let S^1 be the unit circle in $\mathbb{C}$.

Theorem 53. *a.* $\mathrm{H}^\bullet(((\mathbb{CP}^\infty)^m)^n/S_n) \cong \mathrm{Sym}_{A_n}(m)$.

b. $\mathrm{H}^\bullet(((S^1)^m)^n/S_n) \cong \mathrm{Sym}^n(\bigwedge[\theta_1, \dots, \theta_m])$.

c. $\mathrm{H}^\bullet(((\mathbb{CP}^\infty)^m \times (S^1)^k)^n/S_n) \cong \mathrm{Sym}^n(\mathbb{C}[x_1, \dots, x_n] \otimes \bigwedge[\theta_1, \dots, \theta_k])$.

Proof. Recall that $\mathrm{H}^\bullet(\mathbb{CP}^n) = \mathbb{C}[x]/(x^n)$, see [2], which implies that $\mathrm{H}^\bullet(\mathbb{CP}^\infty) = \mathbb{C}[x]$. Thus, by the remarks above

$$\begin{aligned} \mathrm{H}^\bullet(((\mathbb{CP}^\infty)^m)^n/S_n) &\cong ((\mathrm{H}^\bullet(\mathbb{CP}^\infty)^{\otimes m})^{\otimes n})_{S_n} \cong \mathrm{H}^\bullet((\mathbb{C}[x]^{\otimes m})^{\otimes n})_{S_n} \\ &\cong (\mathbb{C}[x_1, \dots, x_m])^{\otimes n}/S_n \cong \mathrm{Sym}_{A_n}(m). \end{aligned}$$

Since $\mathrm{H}^\bullet(S^1) \cong \mathbb{C}[\theta]/\langle\theta^2\rangle$ as graded superalgebras with θ of degree 1, then

$$\mathrm{H}^\bullet(((S^1)^m)^n/S_n) \cong ((\mathrm{H}^\bullet(S^1)^{\otimes m})^{\otimes n})_{S_n} \cong (\bigwedge[\theta_1, \dots, \theta_m]^{\otimes n})_{S_n} \cong \mathrm{Sym}^n(\bigwedge[\theta_1, \dots, \theta_m]).$$

Finally

$$\mathrm{H}^\bullet(((\mathbb{CP}^\infty)^m \times (S^1)^k)^n/S_n) \cong (\mathrm{H}^\bullet(\mathbb{CP}^\infty)^{\otimes m} \otimes \mathrm{H}^\bullet(S^1)^{\otimes k})^{\otimes n}/S_n \cong \mathrm{Sym}^n(\mathbb{C}[x_1, \dots, x_m] \otimes \bigwedge[\theta_1, \dots, \theta_k]).$$

□

Theorem 53 together with the quantizations of $\mathrm{Sym}_{A_n}(2m)$ and of $\mathrm{Sym}^n(\bigwedge[\theta_1, \dots, \theta_m])$ provided in Sections 6.2 and 6.4 respectively, give a quantum product on the cohomology of $((\mathbb{CP}^\infty)^{2m})^n/S_n$ and $((S^1)^m)^n/S_n$ respectively . Notice that the quantum product on $\mathrm{H}^\bullet(((\mathbb{CP}^\infty)^m)^n/S_n)$ is non-commutative and therefore is different to the quantum cohomology product defined for example in [7].

Acknowledgment

We thank Nicolas Andruskiewitsch for his advices and encouragement. We also thank Delia Flores de Chela.

References

[1] J. Alev, T. J. Hodges, and J. D. Velez, *Fixed rings of the Weyl algebra* $A_1(\mathbb{C})$, Journal of algebra **130** (1990), 83–96.

[2] Raoul Bott and Loring W. Tu, *Diferential forms in algebraic topology*, no. 82, Springer-Velarg, New York, 1982.

[3] P. Deligne, P. Etingof, D. Freed, L. Jeffrey, D. Kazhdan, J. Morgan, D. Morrison, and E. Witten, *Quantum fields and strings: A course for mathematicians*, vol. 1, American mathematical society, 1999.

[4] Rafael Díaz and Eddy Pariguan, *Super, quantum and non-commutative species*, Work in progress, 2003.

[5] ———, *Symmetric quantum Weyl algebras*, Annales Mathematiques Blaise Pascal **11** (2004), 155–171.

[6] Vasily Dolgushev, *Covariant and equivariant formality theorems*, math. QA/0307212, 2003.

[7] D. Mc Duff and D. Salamon, *J-holomorphic curves and quantum cohomology*, vol. 6, Univesity lecture series. AMS, Providence, Rhode Island, 1991.

[8] P. Etingof and V. Ginzburg, *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism*, Invent. Math **147** (2002), no. 2, 243–348.

[9] R. Gopakumar, S. Minwalla, and A. Strominger, *Noncommutative Solitons*, J. High Energy Phys. JHEP **05-020** (2000).

[10] J. A. Green, *Polynomial representations of* GL_n, no. 830, Lecture notes in mathematics. Springer-Velarg, New York, 1980.

[11] F. Hirzebruch and T. Höfer, *On the Euler number of an orbifold*, Mathematische Annalen **286** (2002), 255–260.

[12] A. Joyal, *Une théorie combinatoire des séries formelles*, Advances in mathematics (1981), no. 42, 1–82.

[13] ———, *Foncteurs analytiques et espécies de structures*, Combinatoire énumérative. Lecture notes in mathematics, 1234 (1986), 126–159.

[14] M. Kontsevich, *Deformation Quantization of Poisson Manifolds I*, math. q-alg/9709040, 1997.

[15] I. Kriz and J. P. May, *Operads, algebras, modules and motives*, Astérisque (1995), no. 233.

[16] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Mathematical Monographs, New York, 1995.

[17] Emil Martinec and Gregory Moore, *Noncommutative Solitons on Orbifolds*, hep-th/0101199, 2001.

[18] S. A. Merkulov, *The Moyal product in the matrix product*, math-ph/0001039, 2000.

[19] Gian-Carlo Rota, *Gian-Carlo Rota on Combinatorics. Introductory papers and commentaries*, Birkhäuser, Boston, 1995.

[20] R. P. Stanley, *Enumerative combinatorics*, vol. 2, Cambridge University Press, 1999.

[21] F. Vaccarino, *The vector invariants of symmetric groups*, RA/0205233, 2002.

[22] K. H. Wehrhahn, *Combinatorics. An introduction*, vol. 2, Carslaw Publications, Australia, 1990.

Rafael Díaz. Instituto Venezolano de Investigaciones Científicas (IVIC). `radiaz@ivic.ve`
Eddy Pariguan. Universidad Central de Venezuela (UCV). `eddyp@euler.ciens.ucv.ve`